

Frequency and antimicrobial behaviour of thermophilic *Campylobacter* species isolated from Ecuadorian backyard chickens[#]

Frecuencia y comportamiento frente a antimicrobianos de especies termófilas de *Campylobacter* aisladas de pollos de traspatio en Ecuador

S Ochoa^a, R J Simaluiza^a, Z Toledo^a, H Fernández^{b*}

ABSTRACT. The prevalence and antimicrobial behaviour of *Campylobacter jejuni/coli* isolated from backyard chickens from Southern Ecuador were determined. This study revealed that *Campylobacter* sp. frequency was 41.7% with high resistance to tetracycline (94.0%) and ciprofloxacin (88%). *C. jejuni* was the most frequently isolated species (32.5%), followed by *C. coli* (9.2%) being poultry colonization by *Campylobacter* not region dependent. This is the first study on *Campylobacter* antimicrobial resistance in backyard chickens in Southern Ecuador.

Key words: *Campylobacter*, backyard chickens, antimicrobial resistance, Ecuador.

RESUMEN. La prevalencia y el comportamiento antimicrobiano de cepas de *Campylobacter jejuni/coli* aisladas de pollos de traspatio del sur de Ecuador fue determinada. Este estudio revela que la frecuencia de aislamiento de *Campylobacter* sp. fue de 41,7%, observándose una alta resistencia a la tetraciclina (94,0%) y a la ciprofloxacina (88%). *C. jejuni* fue la especie más frecuente (32,5%), seguida de *C. coli* (9,2%), observándose que la colonización de las aves por *Campylobacter* no es dependiente de la región de origen. Este es el primer estudio acerca de la resistencia a los antimicrobianos en *Campylobacter* aislados de pollos de traspatio en el sur de Ecuador.

Palabras clave: *Campylobacter*, pollos de traspatio, resistencia antimicrobiana, Ecuador.

INTRODUCTION

Campylobacter species, particularly *C. jejuni* and *C. coli*, are the most frequent bacterial etiology of human gastroenteritis associated with foodborne diseases worldwide (Humphrey *et al* 2007, Whiley *et al* 2013, WHO 2013).

Acquisition of *Campylobacter* infections occurs through consumption of contaminated food, mainly of poultry origin, water and unpasteurized milk as well as through direct contact with animals and fowls or their faeces (Humphrey *et al* 2007, Fernández 2011, Whiley *et al* 2013, WHO 2013).

In South American and other developing countries, *Campylobacter* infections occur most frequently in children (Fernández 2011). Few data exist regarding the frequency of *Campylobacter* species in children with diarrhoea in Ecuador where these bacteria have been isolated between 10 and 23% (Guderian *et al* 1987, Vasco *et al* 2014).

In high income countries, 58 to 76% of human cases of campylobacteriosis could be attributed to poultry sources (Mullner *et al* 2009, Batz *et al* 2011). However, in developing countries, probably due to epidemiological

data gaps, the possible risk pathways for *Campylobacter* infection via poultry sources are unknown or not well outlined and understood.

Small backyard poultry flocks are popular in Southern Ecuador (Vilcabamba, Loja and Zamora cities) as an economical subsistence way and also as protein source; however there is no epidemiological information about *Campylobacter* in this kind of fowl in this region. For these reasons, the aim of this study was to determine the prevalence and the antimicrobial behaviour of thermophilic *Campylobacter* species isolated from domestic backyard chickens in Southern Ecuador.

MATERIAL AND METHODS

Cloacal swabs were randomly collected from a total of 120 chickens, corresponding 41 to birds from Vilcabamba (4 to 12 birds from 6 backyards), 32 from Loja (5 to 11 birds from 5 backyards) and 47 from Zamora regions (all the birds from the same backyard). The birds were maintained in inadequate environmental cleanliness conditions and in contact with other animal species known as reservoirs for *Campylobacter* like pigs, dogs, cats, sparrows and other free living birds. The cloacal samples collected were seeded into the semisolid TEM transport-enrichment medium described by Fernández (1992) consisting of (formule/L) Brucella broth 28 g, agar 1.5 g, sodium metabisulphite 0.5 g, ferrous sulphate 0.5 g, sodium pyruvate 0.5 g, trimethoprim 10 mg, rifampicin 15 mg, colistin 10,000 IU, amphotericine 10 mg and defibrinated horse blood 30 mL, and transported to the laboratory. Following an enrichment period of 24

Accepted: 25.02.2016.

^aDepartamento de Ciencias de la Salud, Sección Genética Humana, Microbiología y Bioquímica Clínica, Universidad Técnica Particular de Loja, Loja, Ecuador.

^bInstituto de Microbiología Clínica, Facultad de Medicina, Universidad Austral de Chile, Valdivia, Chile.

[#]This work was supported by Project PRO_CCNN_863 UTPL 98739155.

*Corresponding author: H Fernández; hfernand@uach.cl

h at 42 °C under microaerobic conditions obtained using generator envelopes (Oxoid®), aliquots of each sample were seeded onto Butzler medium plates that were incubated for 48 h under the same conditions described above. Suspected colonies were identified morphologically (Gram stain), biochemically (oxidase, catalase, sensitivity to nalidixic acid and cephalotin, hippurate and indoxylacetate hydrolysis) and confirmed by the multiplex PCR test proposed by Yamazaki-Matsune *et al* (2007). In brief, after extraction of genomic DNA, 5 µL of template DNA were added to a mixture containing 2 µM of each primer and 25 µL of 2x Multiplex PCR Master Mix (QIAGEN, USA) adjusting final volume to 50 µL with RNase free H₂O. DNA amplification was performed in thermocycler (Eppendorf) using the following cycling conditions: one cycle of initial denaturation at 95 °C for 15 min, followed by 25 cycles of denaturation at 95 °C for 0.5 min; annealing was done at 58 °C for 1.5 min and extension at 72 °C for 1 min, ending with a final extension time at 72 °C for 7 min. Reaction mixtures were analyzed by gel electrophoresis (3% w/v agarose) stained with ethidium bromide and visualised under UV transilluminator.

Susceptibility to ampicillin, amoxicillin/clavulanic acid, erythromycin, tetracycline, ciprofloxacin and gentamycin was determined by the disk diffusion method following the 2014 recommendations of The European Committee on Antimicrobial Susceptibility Testing-EUCAST and the Committee for the Antibiogram of the French Society of Microbiology (Société Française de Microbiologie 2014).

Data were analysed by means of the R Statistical Computing Package (Vienna, Austria, 2005) using Fisher's exact two-tailed test with a significance level of 5%.

RESULTS AND DISCUSSION

The results obtained are shown in table 1. From the 120 chickens studied, 50 (41.7%) harbored thermophilic *Campylobacter* species being *C. jejuni* the most frequently isolated (39/120, 32.5%) followed by *C. coli* (11/120, 9.2%).

With regard to the frequency for each location, the lowest isolation frequency was obtained from chickens samples collected in Loja city (37.5%). However, the

observed differences among the three groups of chickens were not statistically significant ($P > 0.005$).

Table 2 shows that the isolated strains showed very high resistant frequencies to tetracycline (94.0%; *C. jejuni* 94.9%, *C. coli* 91.0%) and ciprofloxacin (88.0%; *C. jejuni* 84.6%, *C. coli* 100%). Resistance to ampicillin was observed in 12 strains (24.0%; *C. jejuni* 15.4%, *C. coli* 54.5%). All the strains were susceptible to erythromycin, gentamycin and amoxicillin/clavulanic acid. Multi-resistance (resistance to three antimicrobials) was found in 12 (24%) strains, the same that were resistant to ampicillin. Multi-resistance was most frequent amongst *C. coli* (7/11 strains, 63.6%) than in *C. jejuni* (4/11 strains, 36.4%). All the ampicillin resistant strains were susceptible to amoxicillin/clavulanic acid.

The prevalence of the thermotolerant species of *Campylobacter* found in this study (41.7%) was higher than the prevalence reported in Southern Chile (25.7%) by Fernández and Torres (2000) but lower to that informed in New Zealand, where 86% of flocks tested positive with a *Campylobacter* isolation rate of 57.7% (Anderson *et al* 2012). However, in agreement with the results of these authors, our isolation rate of *C. jejuni* was higher than the rate obtained for *C. coli*. This was observed in the three groups of chicken studied and seems to be a fact regularly observed in different kinds of samples, at least in Latin American countries (Fernández 2011). Similar results were recently obtained studying chicken livers for human consumption in Southern Ecuador (Simaluiza *et al* 2015).

Surprisingly, we found a remarkable high resistance rate to tetracycline (94.0%) and to ciprofloxacin (88.0%). Previously, high resistance levels were also found in *Campylobacter* strains isolated from chicken livers disposed for human consumption in this same region (Simaluiza *et al* 2015). However we could not obtain information from the owners about the use of antibiotics, we believe that as occurs in other places of the developing world, this probably reflects an overuse of these antibiotics in backyard chickens in this region, something that could represent an epidemiological problem with implications for treatment in human beings (WHO 2013, Whitehead and Roberts 2014). This is particularly important in backyard chickens because i) exposure of children to infected backyard poultry may represent another route of transmission for

Table 1. Occurrence of *Campylobacter* species among backyard chickens from three locations of Southern Ecuador.
Prevalencia de especies de *Campylobacter* en pollos de traspatio de tres localidades del sur de Ecuador.

Location of Origin	Number of samples	<i>Campylobacter</i> positive N°%	<i>C. jejuni</i> N°%	<i>C. coli</i> N°%
Vilcabamba	41	17/41* 41.5	13/17 76.5	4/17 23.5
Loja	32	12/32* 37.5	10/12 83.3	2/12 16.7
Zamora	47	21/47* 44.7	16/21 76.2	5/21 23.8
TOTAL	120	50/120 41.7	39/50 78.0	11/50 22.0

* $P > 0.005$.

Table 2. Antimicrobial behavior of *Campylobacter* species isolated from backyard chickens from three locations of Southern Ecuador. Comportamiento frente a antimicrobianos de *Campylobacter* aislados de pollos de traspatio de tres localidades del sur de Ecuador.

Location of Origin (n)	Amp S R	Amo-Clav S R	Genta S R	Tet S R	Ery S R	Cip S R
Vilcabamba						
<i>C. jejuni</i> (13)	11/13 2/13 (84.6%) (15.4%)	13/13 0 (100%) (0%)	13/13 0 (100%) (0%)	0 13/13 (0%) (100%)	13/13 0 (100%) (0%)	0 13/13 (0%) (100%)
<i>C. coli</i> (4)	1/4 3/4 (25%) (75%)	4/4 0 (100%) (0%)	4/4 0 (100%) (0%)	0 4/4 (0%) (100%)	4/4 0 (100%) (0%)	0 4/4 (0%) (100%)
Loja						
<i>C. jejuni</i> (10)	6/10 4/10 (60%) (40%)	10/10 0 (100%) (0%)	10/10 0 (100%) (0%)	0 10/10 (0%) (100%)	10/10 0 (100%) (0%)	0 10/10 (0%) (100%)
<i>C. coli</i> (2)	1/2 1/2 (50%) (50%)	2/2 0 (100%) (0%)	2/2 0 (100%) (0%)	0 2/2 (0%) (100%)	2/2 0 (100%) (0%)	0 2/2 (0%) (100%)
Zamora						
<i>C. jejuni</i> (16)	16/16 0 (100%) (0%)	16/16 0 (100%) (0%)	16/16 0 (100%) (0%)	2/16 14/16 (12.5%) (87.5%)	16/16 0 (100%) (0%)	6/16 10/16 (37.5%) (62.5%)
<i>C. coli</i> (5)	3/5 2/5 (60%) (40%)	5/5 0 (100%) (0%)	5/5 0 (100%) (0%)	1/5 4/5 (20%) (80%)	5/5 0 (100%) (0%)	0 5/5 (0%) (100%)
TOTAL						
<i>C. jejuni</i> (39)	33/39 6/39 (84.6%) (15.4%)	39/39 0 (100%) (0%)	39/39 0 (100%) (0%)	2/39 37/39 (5.1%) (94.9%)	39/39 0 (100%) (0%)	6/39 33/39 (15.4%) (84.6%)
<i>C. coli</i> (11)	5/11 6/11 (45.5%) (54.5%)	11/11 0 (100%) (0%)	11/11 0 (100%) (0%)	1/11 10/11 (9%) (91%)	11/11 0 (100%) (0%)	0 11/11 (0%) (100%)

S = susceptible, R = resistant, Amp = ampicillin, Amo-Clav = amoxicillin-clavulanic acid, Genta = gentamycin, Tet = tetracycline, Ery = erythromycin, Cip = ciprofloxacin.

Campylobacter infection (Anderson *et al* 2012, Whitehead and Roberts 2014, El-Tras *et al* 2015), on which there are not sufficient epidemiological studies and ii) there is a lack of good biosecurity and good management practices applicable to reduce the risk of infection and antimicrobial resistance levels in backyard chickens (Whitehead and Roberts 2014, El-Tras *et al* 2015).

All the strains were susceptible to gentamycin, erythromycin and amoxicillin-clavulanic acid but 24.0% were resistant to ampicillin. Resistance to ampicillin and other beta-lactam antibiotics has been found in *Campylobacter* strains isolated from humans, poultry and poultry products. Since the ampicillin resistant strains were amoxicillin/clavulanic acid susceptible, we can infer that the resistance to ampicillin could be mediated by the production of a beta lactamase (Simaluiza *et al* 2015).

Multi-resistance was found in 24% of the strains being most frequent amongst *C. coli* than in *C. jejuni* strains. It is known that *C. coli* are usually more resistant than *C. jejuni* strains. This was also found in *C. coli* isolated from chicken livers in Southern Ecuador (Simaluiza *et al* 2015).

According to the results, antimicrobial resistance to tetracycline and ciprofloxacin in *Campylobacter* strains is

a serious epidemiological problem in Southern Ecuador, whereas ampicillin resistance may be an emerging problem which could require the establishment of a laboratory surveillance program to assess its real significance. On the other hand, good biosecurity and management practices are needed to be applied to prevent the transmission of *Campylobacter* from poultry to children and to control antimicrobial resistance spread. Among good biosecurity and management practices are: the use of clean water, avoiding cross contamination through contact with other animals in the same farmyard as well as the implementation of basic sanitation measures in order to obtain better environmental cleanliness.

As far as we know, this is the first attempt to put in evidence the occurrence of *Campylobacter* species in backyard chickens in Southern Ecuador, as well their antimicrobial susceptibility and resistance profiles.

ACKNOWLEDGEMENTS

This work was supported by Project PRO_CCNN_863 UTPL 98739155. H. Fernández was supported by the Prometeo Project of the Secretariat for Higher Education, Science, Technology and Innovation of the Republic of Ecuador.

REFERENCES

- Anderson J, BJ Horn, BJ Gilpin. 2012. The Prevalence and genetic diversity of *Campylobacter* spp. in domestic 'backyard' poultry in Canterbury, New Zealand. *Zoonoses Public Health* 59, 52-60.
- Batz B, S Hoffmann, G Morris. 2011. Ranking the Risks: The 10 pathogen-food combinations with the greatest burden on public health. University of Florida, Emerging Pathogens Institute. Gainesville, Florida, USA, 1-70.
- El-Tras WF, HR Holt, AA Tayel, NN El-Kady. 2015. *Campylobacter* infections in children exposed to infected backyard poultry in Egypt. *Epidemiol Infect* 143, 308-315.
- Fernández H. 1992. Increase of *Campylobacter* isolation rates using an enrichment medium. *Rev Microbiol São Paulo* 23, 5-7.
- Fernández H, N Torres. 2000. Occurrence of *Campylobacter jejuni* and *Campylobacter coli* in three groups of hens of different geographic origin in Southern Chile. *Arch Med Vet* 32, 241-244.
- Fernández H. 2011. *Campylobacter* and campylobacteriosis: a view from South America. *Rev Peruana Med Exp Salud Pública* 28, 121-127.
- Guderian R, G Ordóñez, R Bossano. 1987. Acute diarrhea associated with *Campylobacter* and other pathogens in Quito, Ecuador. *B Ofic Sanit Panam* 102, 333-339.
- Humphrey T, S O'Brien, M Madsen. 2007. Campylobacters as zoonotic pathogens: A food production perspective. *Int J Food Microbiol* 117, 237-257.
- Mullner P, SEF Spencer, D Wilson, G Jones, AD Noble, AC Midwinter, JM Collins-Emerson, P Carter, S Hathaway, NP French. 2009. Assigning the source of human campylobacteriosis in New Zealand: a comparative genetic and epidemiological approach. *Infect Genet Evol* 9, 1311-1319.
- Simaluiza RJ, Z Toledo, S Ochoa, H Fernández. 2015. Resistance of *Campylobacter jejuni* and *Campylobacter coli* in chicken livers used for human consumption in Ecuador. *J Anim Vet Adv* 14, 6-9.
- Société Française de Microbiologie. 2014. *Comité de l'Antibiogramme-Recommandations* 2014. Société Française de Microbiologie, Paris, France, Pp 1-115.
- Vasco G, G Trueba, R Atherton, M Calvopiña, W Cevallos, T Andrade, M Eguiguren, JN Eisenberg. 2014. Identifying etiological agents causing diarrhea in low income Ecuadorian communities. *Am J Trop Med Hyg* 91, 563-569.
- Whiley H, B van den Akker, S Giglio, R. Bentham 2013. The role of environmental reservoirs in human campylobacteriosis. *Int J of Environ Res Public Hlth* 10, 5886-5907.
- Whitehead ML, V Roberts. 2014. Backyard poultry: legislation, zoonoses and disease prevention. *J Small Anim Pract* 55, 487-496.
- WHO, World Health Organization. 2013. The global view of campylobacteriosis. Report of an expert consultation. *WHO Document Production Services*, Geneva, Switzerland.
- Yamazaki-Matsune W, M Taguchi, K Seto, R Kawahara, K Kawatsu, Y Kumeda, M Kitazato, M Nukina, N Misawa, T Tsukamoto. 2007. Development of a multiplex PCR assay for identification of *Campylobacter coli*, *Campylobacter fetus*, *Campylobacter hyointestinalis* subsp. *hyointestinalis*, *Campylobacter jejuni*, *Campylobacter lari* and *Campylobacter upsaliensis*. *J Med Microbiol* 56, 1467-1473.