

First record of the eucalypt gall-wasp *Leptocybe invasa* (Hymenoptera: Eulophidae) from Uruguay

Primer reporte de la avispa agalladora del eucalipto *Leptocybe invasa* (Hymenoptera: Eulophidae) para Uruguay

Carolina Jorge^{**}, Gonzalo Martínez^b, Demian Gómez^b, Martín Bollazzi^c

*Autor de Correspondencia: ^aUniversidad de la República, CENUR Noreste sede Tacuarembó, Instituto Superior de Estudios Forestales, Ruta 5, km 387, Tacuarembó, Uruguay, tel.: 46333485 int. 262, carolina.jorge@cut.edu.uy

^bINIA Tacuarembó, Programa Nacional de Producción Forestal, Laboratorio de Entomología, Tacuarembó, Uruguay.

^cUniversidad de la República, Facultad de Agronomía, Departamento de Protección Vegetal, Unidad de Entomología, Montevideo, Uruguay.

SUMMARY

Leptocybe invasa is a recently introduced forest pest, responsible for significant economic loss in *Eucalyptus* spp. plantations worldwide. The objective of this work is to report its presence in Uruguay, providing data of its distribution and biology. A survey for galls was conducted from 2013 to 2015. The presence of galls was confirmed in 15 out of 35 surveyed points and four out of 157 traps. *Eucalyptus benthamii* was recorded as a new host. Three periods of emergence were observed. Life cycle was completed only in *E. tereticornis*, *E. camaldulensis* and *E. grandis*.

Key words: gall-insects, *Eucalyptus*, forestry pests.

RESUMEN

Leptocybe invasa es una plaga forestal de reciente introducción, responsable de importantes pérdidas económicas en plantaciones de *Eucalyptus* spp. a nivel mundial. El objetivo de este trabajo fue reportar su presencia para Uruguay, aportar datos de distribución y biología. Se observó la presencia de agallas de 2013 a 2015, confirmada en 15 de 35 puntos muestreados y en cuatro de 157 trampas. *Eucalyptus benthamii* se registra como nuevo hospedero. Fueron observados tres periodos de emergencia de adultos. El ciclo de vida fue completado solamente en *E. tereticornis*, *E. camaldulensis* y *E. grandis*.

Palabras clave: insectos agallícolas, *Eucalyptus*, plagas forestales.

INTRODUCTION

Commercial forestry with *Eucalyptus* species reaches 80 % of the forested area in Uruguay, leading, in 2013, to more than 590 million dollars in export goods (DIEA 2014). *Eucalyptus globulus* Labill and *E. grandis* W. Hill ex Maiden are the most planted species, while red gum trees (*E. tereticornis* Sm. and *E. camaldulensis* Dehnh.) have been extensively used as shade and shelter forest in cattle breeding. Paine *et al.* (2011) suggest a correlation between the increase of global area planted with *Eucalyptus* spp. and the arrival of new insects and diseases. This phenomenon has also been observed in Uruguay, particularly during the last decade (Martínez 2010).

The last wave of world invasive forest pests has been characterized by small, cryptic insects, notably gall forming species (Paine *et al.* 2011). The *Eucalyptus* gall-wasps *Ophelimus maskelli* Ashmead 1900, *Selitrichodes*

globulus La Salle and Gates 2009 and *Leptocybe invasa* Fischer and La Salle 2004 are well known because of their economic importance, though the latter is particularly relevant due to the aggressiveness of the attack and its rapid dispersion (Paine *et al.* 2011).

The Australian gall wasp *L. invasa* was first recorded from South America in 2008; in Brazil in 2008, Argentina in 2010 and Chile in 2014 (Wylie and Speight 2012, SAG 2014). This insect has become a relevant pest for eucalypt forestry worldwide. Life cycle occurs almost entirely inside the gall. In temperate regions, the developmental cycle lasts 130 days on average leading to the occurrence of 2-3 generations annually (Mendel *et al.* 2004, Aquino *et al.* 2011), reaching to 5-6 generations in tropical areas (Zhu *et al.* 2012). Thelytokous parthenogenesis is the most common reproductive mechanism in *L. invasa* (Mendel *et al.* 2004). The objective of this work is to report the presence of *L. invasa* in Uruguay and the first mention of *Eucalyptus benthamii* as its host.

METHODS

Occurrence of galls on eucalypt was recorded for the first time in April 2011 in Montevideo during an expert visit (34°50'10.96" S; 56°13'11.77" W). From August 2013 to August 2015 we conducted a survey of galls on eucalypt trees in plantations and shade and shelter forests. Thirty-five sites were selected following two criteria: (1) presence of *Eucalyptus* species susceptible to *L. invasa* and (2) proximity to wood transport routes. All sites were inspected monthly to observe gall development. Adult wasps for identification were obtained from two sites located on *E. tereticornis* x *E. camaldulensis* hybrids and *E. grandis* at INIA Experimental Station (Tacuarembó). Additional points of occurrence were obtained from the review of 157 yellow sticky traps belonging to the National Monitoring System (Martínez *et al.* 2014).

The observation of specimens was made under a stereoscopic microscope Zeiss Stemi DV4 and by Scanning Electron Microscope (SEM) JEOL 5900 Low Vacuum. We followed Mendel *et al.* (2004) to determine the adult wasps and to describe gall morphology and we compared the individuals against identified material kept in the Forest Entomological Collection of INIA in Tacuarembó, Uruguay.

RESULTS

Gall and adult sampling. Gall occurrence was confirmed on 15 out of 35 surveyed points (figure 1). Emergency was confirmed on *E. tereticornis* x *E. camaldulensis* and *E. grandis* in three periods: August-September, mid-December and March-April. Only four out of 157 sticky traps checked contained adult wasp of *L. invasa* within the same three periods observed in the field survey (figure 1).

Material examined. Uruguay, Tacuarembó: INIA Experimental Station (31°44'23.84" S; 55°58'44.74" W) ex *Eucalyptus tereticornis* Sm. (three female adults).

Leptocybe invasa is a small wasp (1.1-1.4 mm) brownish in color with blue green metallic shine on the thorax (figure 2A). Specimens were ascribed to *L. invasa* based on the antennal morphology, thoracic morphology and wing venation (figure 2B-D). Detailed diagnosis of adult female was provided by Mendel *et al.* (2004). Scape longer than the *pedicellum* with broadened middle section. *Flagellum* with six segments separated by four *annelli*, the basal segments constitute the *funiculum* and the three apical segments form the antennal mass (figure 2B). Short prothorax; well-developed mesothorax; *scutellum* divided into three




Figure 1. Occurrence of *L. invasa* in Uruguay. On yellow sticky cards: grey dots = presence; white dots = absence; black dots = presence of galls on survey points.

Presencia de *L. invasa* en Uruguay. En trampas amarillas: puntos grises = presencia; puntos blancos = ausencia; puntos negros = presencia de agallas en puntos inspeccionados.

separated zones by sublateral lines (figure 2C). Hyaline wings completely covered with setae, slight wing venation (figure 2D). Post marginal vein is shorter than stigmal vein. Submarginal vein presents 2-5 setae, visible on dorsal view. Identified material was deposited in the Forestry Entomological Collection of INIA in Tacuarembó, Uruguay.

Galls of *L. invasa* are always located in the central rib or in leaf petioles (figure 3A). Early attack is observed in

the form of rectilinear wounds on the epidermis, which are inflicted by females during oviposition (figure 3B). Induced galls can cause deformation on leaves and apical shoots (figure 3C). Severe attacks can produce tissue necrosis and defoliation (figure 3D).

Table 1 reviews eucalypt species reported as susceptible to *L. invasa*. We observed for the first time the occurrence of galls on *E. benthamii*. Additionally, we observed




Figure 2. Adult female of *Leptocybe invasa*: A) Lateral view, body length 1.2 mm. B) Antenna. C) Thorax. D) Wings.
Hembra adulta de *L. invasa*: A) Vista lateral, longitud del cuerpo 1,2 mm. B) Antena. C) Tórax. D) Alas.




Figure 3. A) Shape and typical position of galls of *Leptocybe invasa* in *Eucalyptus tereticornis* x *E. camaldulensis*. B) Oviposition marks in *E. globulus*. C) Anomalous development on young plant in *E. grandis*. D) Necrosis and death of leaves and stems as a consequence of the attack by *L. invasa* in *E. globulus*.

A) Forma y posición típica de agallas de *L. invasa* en *E. tereticornis* x *E. camaldulensis*. B) Marcas de oviposición en *E. globulus*. C) Desarrollo anómalo en plantas jóvenes de *E. grandis*. D) Necrosis y muerte de hojas y ramas a consecuencia del ataque de *L. invasa* en *E. globulus*.

Table 1. Reported hosts for *Leptocybe invasa*. N.S.: Not specified.

Hospederos reportados para *Leptocybe invasa*. N.S.: No especificado.

Species	Countries/Sources
Subgenus: <i>Idiogenes</i>	
<i>E. cloeziana</i> F. Muell	N.S. (Nadel and Slippers 2011)
Subgenus: <i>Monocalyptus</i>	
<i>E. pilularis</i> Sm.	Vietnam (Quang Thu <i>et al.</i> 2009), N.S. (Nadel and Slippers 2011)
Subgenus: <i>Nothocalyptus</i>	
<i>E. microcorys</i> Muell.	Vietnam (Quang Thu <i>et al.</i> 2009), N.S. (Nadel and Slippers 2011), China (Zhu <i>et al.</i> 2012)
Subgenus: <i>Symphomyrtus</i>	
<i>Eucalyptus alba</i> Reinw. ex Blume	N.S (Nadel and Slippers 2011)
<i>E. botryoides</i> Sm.	Israel (Mendel <i>et al.</i> 2004), N.S. (Nadel and Slippers 2011)
<i>E. bridgesiana</i> Baker	Israel (Mendel <i>et al.</i> 2004), N.S. (Nadel and Slippers 2011), N.S. (FAO 2012)
<i>E. camaldulensis</i> Dehnh.	Israel (Mendel <i>et al.</i> 2004), Uganda (Nyeko <i>et al.</i> 2010), Vietnam (Quang Thu <i>et al.</i> 2009), Argentina (Aquino <i>et al.</i> 2011, Nadel and Slippers 2011), N.S (FAO 2012), China (Zhu <i>et al.</i> 2012, Luo <i>et al.</i> 2014), Brazil (Cegatta and Villegas 2013), Mozambique (Chirinzane <i>et al.</i> 2014) Chile (SAG 2014)
<i>E. cinerea</i> F. Muell. ex Benth.	N.S (FAO 2012)
<i>E. cladocalyx</i> Muell.	N.S (Nadel and Slippers 2011)
<i>E. coolabah</i> Blakely et Jacobs	Vietnam (Quang Thu <i>et al.</i> 2009)
<i>E. dunnii</i> Maiden	Kenya (Nyeko <i>et al.</i> 2010), N.S. (Nadel and Slippers 2011), N.S. (FAO 2012), China (Zhu <i>et al.</i> 2012)
<i>E. excerta</i> Muell.	China (Zhu <i>et al.</i> 2012)
<i>E. globulus</i> Labill.	Israel (Mendel <i>et al.</i> 2004), Uganda (Nyeko <i>et al.</i> 2009), N.S (Nadel and Slippers 2011), N.S. (FAO 2012) Chile (SAG 2014)
<i>E. gomphocephala</i> D.C.	N.S. (Nadel and Slippers(2011)
<i>E. grandis</i> Hill ex Maiden	Israel (Mendel <i>et al.</i> 2004), Vietnam (Quang Thu <i>et al.</i> 2009), Kenya and Uganda (Nyeko <i>et al.</i> 2010), N.S. (Nadel and Slippers 2011), N.S (FAO 2012)
<i>E. gunnii</i> Hook. F.	Israel (Mendel <i>et al.</i> 2004)
<i>E. leucoxyton</i> Muell.	N.S. (Nadel and Slippers 2011)
<i>E. maidenii</i> Muell.	Uganda (Nyeko <i>et al.</i> 2009), Vietnam (Quang Thu <i>et al.</i> 2009), N.S. (Nadel and Slippers 2011), N.S. (FAO 2012)
<i>E. moluccana</i> Roxb.	Vietnam (Quang Thu <i>et al.</i> 2009)
<i>E. nicholii</i> Maiden et Blakely	N.S. (FAO 2012)
<i>E. nitens</i> Deane et Maiden	N.S. (Nadel and Slippers 2011)
<i>E. pellita</i> Muell.	Vietnam (Quang Thu <i>et al.</i> 2009), N.S. (Nadel and Slippers 2011)
<i>E. propinqua</i> Deane et Maiden	China (Zhu <i>et al.</i> 2012)
<i>E. pulverulenta</i> Sims	N.S. (FAO 2012)
<i>E. robusta</i> Sm.	Israel (Mendel <i>et al.</i> 2004), Uganda (Nyeko <i>et al.</i> 2010), Vietnam (Quang Thu <i>et al.</i> 2009), N.S. (Nadel and Slippers 2011), N.S. (FAO 2012), China (Zhu <i>et al.</i> 2012)
<i>E. rudis</i> Sm.	N.S. (FAO 2012)
<i>E. saligna</i> Sm.	Israel (Mendel <i>et al.</i> 2004), Vietnam (Quang Thu <i>et al.</i> 2009), Kenya and Uganda (Nyeko <i>et al.</i> 2010), N.S. (Nadel and Slippers 2011), China (Zhu <i>et al.</i> 2012), Mozambique (Chirinzane <i>et al.</i> 2014)
<i>E. sideroxyton</i> Cunn.	N.S (Nadel and Slippers (2011))
<i>E. smithii</i> Baker	Vietnam (Quang Thu <i>et al.</i> (2009)), N.S (Nadel and Slippers (2011))
<i>E. tereticornis</i> Sm.	Israel (Mendel <i>et al.</i> (2004)), Vietnam (Quang Thu <i>et al.</i> (2009)), Kenya (Nyeko <i>et al.</i> (2010)), N.S (Nadel and Slippers (2011)), China (Zhu <i>et al.</i> (2012), Luo <i>et al.</i> (2014))
<i>E. urophylla</i> Blake	Vietnam (Quang Thu <i>et al.</i> (2009)), Kenya (Nyeko <i>et al.</i> (2010)), N.S (Nadel and Slippers (2011)), N.S (FAO (2012)), China (Zhu <i>et al.</i> (2012))
<i>E. viminalis</i> Labill.	Israel (Mendel <i>et al.</i> (2004)), N.S (Nadel and Slippers (2011))

galls on red gums (*E. tereticornis*, *E. camaldulensis* and hybrids of these two species), *E. dunnii*, *E. grandis* and *E. globulus*. We only observed emergency holes on red gums and *E. grandis* in different growing stages in the field, we did not inspect greenhouses or nurseries.

DISCUSSION

We observed galls on six species already reported as susceptible, with the addition of *E. benthamii*. Emergence period observed in our study suggests that the duration of the life cycle in Uruguay could be on average 120 days, similar to other temperate regions (Aquino *et al.* 2011). The presence of emergency holes solely on *E. grandis* and red gums suggests that not all species are equally suitable to complete the cycle. However, from the point of view of forestry, gall formation *per se* may hamper the development of the tree stands implying additional costs for tree production.

We assume that *L. invasa* colonized Uruguay from Argentina. Data suggest that, as first reports were closer to the Uruguayan border (provinces of Buenos Aires and Entre Rios) than to Brazil (Bahia and São Paulo States (Costa *et al.* 2008), and no reports have been made yet by foresters from the East region of Uruguay.

We collected additional individuals from a network of yellow sticky traps already established for monitoring other forestry pests. Although sticky traps have been used in other countries for collecting *L. invasa* (Kumari *et al.* 2010, Zhu *et al.* 2012), identification of the captured micro-hymenopteran insects has proven a difficult task due to the conservation status of the specimens.

The report of *L. invasa* in the country poses a threat to Uruguayan commercial forestry, hence research should focus on the development of management strategies for this pest. Chemical control has been tested in Brazil and India (Kumari *et al.* 2010, Cegatta and Villegas 2013), although more than 80 % of the commercial stands are certified under ISO or FSC standards in Uruguay, turning this option unlikely, or at least restricted to nurseries. Alternatively, plant breeding seems to be a promising long term control tool for this pest (Kulkarni 2010, Dittrich-Schröder *et al.* 2012, Zhu *et al.* 2012). Since we observed variation in susceptibility to *L. invasa* among *Eucalyptus* species and hybrids, future studies should focus on screening resistance to attack from *L. invasa* in national genetic stock. Biological control programs are currently being implemented in Israel (Kim *et al.* 2008), South Africa (Kelly *et al.* 2012, Dittrich-Schröder *et al.* 2014), Chile (SAG 2014) and Brazil (de Sá *et al.* 2015). National studies may focus on the identification of the most reliable biocontrol candidates for our ecological context.

ACKNOWLEDGMENTS

The authors want to thank the forestry companies UPM-Forestal Oriental S.A. and Weyerhaeuser Produc-

tos S.A. for the access to their plantations. GMC wants to thank Sociedad de Productores Forestales of Uruguay (SPF) for granting the access to the yellow sticky cards of the National Monitoring Scheme.

REFERENCES

- Aquino AD, EN Botto, MS Loiacono, P Pathauer. 2011. Avispa de la agalla del eucalipto, *Leptocybe invasa* Fisher and La Salle (Hymenoptera: Eulophidae: Tetrastichinae), en Argentina. *Revista INTA* 37:159-164.
- Cegatta IR, C Villegas. 2013. Eficiência de dois inseticidas sistêmicos no controle de *Leptocybe invasa* em mudas de *Eucalyptus camaldulensis*. *Revista do Instituto Florestal* 25:215-221.
- Chirinzane CJ, MD de Souza, NJ Souza, EH Rezende, T Alves, RR Bandeira, AM Fernandes. 2014. First report of *Leptocybe invasa* Fisher and La Salle (*sic*) (Hymenoptera: Eulophidae) in Mozambique. *African Journal of Agricultural Research* 9(49):3555-3558. DOI: 10.5897/AJAR2013.8109.
- Costa VA, E Berti Filho, CF Wilcken, JL Stape, J La Salle and LD Teixeira. 2008. *Eucalyptus* gall wasp, *Leptocybe invasa* Fisher et La Salle (Hymenoptera: Eulophidae) in Brazil: new forest pest reaches the New World. *Revista de Agricultura* 83(2):136-139.
- DIEA (Dirección de Estadísticas Agropecuaria, UY). 2014. Anuario Estadístico 2014. DIEA - MGAP. Accessed Jul. 1st, 2015. Available at: <http://www.mgap.gub.uy/portal/page.aspx?2,diea,diea-anuario-2014,O,es,0,%20>.
- Dittrich-Schröder G, M Harney, S Naser, T Joffe, S Bush, BP Hurley, MJ Wingfield, B Slippers. 2014. Biology and host preference of *Selitrichodes neseri*: A potential biological control agent of the *Eucalyptus* gall wasp, *Leptocybe invasa*. *Biological Control* 78:33-41. DOI: 10.1016/j.biocontrol.2014.07.004
- Dittrich-Schröder G, MJ Wingfield, BP Hurley, B Slippers. 2012. Diversity in *Eucalyptus* susceptibility to the gall-forming wasp *Leptocybe invasa*. *Agricultural and Forest Entomology* 14(4):419-427. DOI: 10.1111/j.1461-9563.2012.00583.x
- FAO (Food and Agriculture Organisation of the United Nation, IT). 2012. *Leptocybe invasa* Fisher and La Salle. FAO - Forestry Department. Accessed Jul. 20th, 2015. Available at: <http://www.fao.org/forestry/13569-05912e0e2fe9054c3e-d4904ae597e3310.pdf>.
- Kelly J, J La Salle, M Harney, G Dittrich-Schröder, BP Hurley. 2012. *Selitrichodes neseri* n. sp, a new parasitoid of the eucalyptus gall wasp *Leptocybe invasa* Fisher and La Salle (Hymenoptera: Eulophidae: Tetrastichinae). *Zootaxa* 3333:50-57.
- Kim IK, Z Mendel, A Protasov, D Blumberg, J La Salle. 2008. Taxonomy, biology, and efficacy of two Australian parasitoids of the eucalyptus gall wasp, *Leptocybe invasa* Fisher and La Salle (Hymenoptera: Eulophidae: Tetrastichinae). *Zootaxa* 1910:1-20.
- Kulkarni HD. 2010. Screening eucalyptus clones against *Leptocybe invasa* Fisher and La Salle (Hymenoptera: Eulophidae). *Karnataka Journal of Agricultural Sciences* 23(1):87-90.
- Kumari NK, HD Kulkarni, KB Goud. 2010. Biology of eucalyptus gall wasp, *Leptocybe invasa* Fisher and La Salle (Hymenoptera: Eulophidae). *Karnataka Journal of Agricultural*

- Sciences* 23(1):211-212.
- Luo J, R Arnold, W Lu, Y Lin. 2014. Genetic variation in *Eucalyptus camaldulensis* and *E. tereticornis* for early growth and susceptibility to the gall wasp *Leptocybe invasa* in China. *Euphytica* 196:397-411. DOI: 10.1007/s10681-013-1042-8.
- Martínez G. 2010. Insectos plaga en plantaciones jóvenes de eucalipto: hacia un modelo. *INIA. Serie Actividades de Difusión* 629:9-24.
- Martínez G, D Gómez, GS Taylor. 2014. First record of the Australian psyllid *Blastopsylla occidentalis* Taylor (Hemiptera, Psylloidea) from Uruguay. *Transactions of the Royal Society of South Australia* 138(2):231-236.
- Mendel Z, A Protasov, N Fisher, J La Salle. 2004. Taxonomy and biology of *Leptocybe invasa* gen. et sp. n. (Hymenoptera: Eulophidae), an invasive gall inducer on *Eucalyptus*. *Australian Journal of Entomology* 43:101-113.
- Nadel RL, B Slippers. 2011. *Leptocybe invasa*, the blue gum chalcid wasp. ICFR, South Africa. Accessed Dec. 20th, 2014. Available at: <http://www.forestry.co.za/uploads/File/home/notices/2011/ICFR%20IS01-2011gallwasp.pdf>.
- Nyeko P, KE Mutitu, BO Otieno, GN Ngae, RK Day. 2010. Variations in *Leptocybe invasa* (Hymenoptera: Eulophidae) population intensity and infestation on eucalyptus germplasms in Uganda and Kenya. *International Journal of Pest Management* 56(2): 137-144
- Nyeko P, EK Mutitu, RK Day. 2009. *Eucalyptus* infestation by *Leptocybe invasa* in Uganda. *African Journal of Ecology* 47(3):299-307.
- Paine TD, MJ Steinbauer, SA Lawson. 2011. Native and Exotic Pests of *Eucalyptus*: A Worldwide Perspective. *Annual Review of Entomology* 56:181-201. DOI: 10.1146/annurev-ento-120709-144817.
- Quang Thu P, B Dell, TI Burgess. 2009. Susceptibility of 18 eucalypt species to the gall wasp *Leptocybe invasa* in the nursery and young plantations in Vietnam. *ScienceAsia* 35(2):113-117. DOI: 10.2306/scienceasia1513-1874.2009.35.113.
- Sá LAN de, AR de Souza, LR Junqueira, MC Candelária, VA Costa, LR Barbosa, JC Zanuncio, CF Wilcken. 2015. Introdução do parasitoide *Selitrichodes neseri* (Hymenoptera: Eulophidae) para o controle da vespa-da-galha do eucalipto *Leptocybe invasa* (Hymenoptera: Eulophidae). Resumos do 14 Simpósio de controle biológico. Teresópolis, Brasil.
- SAG (Servicio Agrícola y Ganadero, CL). 2014. *Leptocybe invasa* Fischer et Lasalle (Hymenoptera: Eulophidae) Microavispa formadora de agallas en eucalipto. Servicio Agrícola y Ganadero. Accessed Jul. 22nd, 2015. Available at: <http://www.sag.cl/ambitos-de-accion/leptocybe-invasa-fischer-lasalle-hymenoptera-eulophidae-microavispa-formadora-de>
- Wylie FR, MR Speight. 2012. Insect pests in tropical forestry. 2nd Edition. Oxfordshire, United Kingdom. CABI. 365 p.
- Zhu F, S Ren, B Qiu, Z Huang, Z Peng. 2012. The Abundance and Population Dynamics of *Leptocybe invasa* (Hymenoptera: Eulophidae) Galls on *Eucalyptus* spp. in China. *Journal of Integrative Agriculture* 11(12):2116-2123. DOI: 10.1016/S2095-3119(12)60470-5.

Recibido: 14.10.15
Aceptado: 29.04.16