



PALEOEPIDEMIOLOGY OF INTESTINAL PARASITES AND LICE IN PRE-COLUMBIAN SOUTH AMERICA*

PALEOEPIDEMIOLOGÍA DE PARÁSITOS INTESTINALES Y PIOJOS EN SUDAMÉRICA PRECOLOMBINA

Adauto Araújo¹, Karl Reinhard², Daniela Leles¹, Luciana Sianto¹, Alena Ñiguez³,
Martín Fugassa⁴, Bernardo Arriaza^{5,6,7}, Nancy Orellana^{6,8,9}, and Luiz Fernando Ferreira¹

Some human parasites originated in prehomínid ancestors in Africa. Nematode species, such as *Enterobius vermicularis* (pinworm), hookworms and *Trichuris trichiura* are shared by humans and other close phylogenetic primates (*Pan* and *Gorilla*), showing that they infected a common ancestor to this group. When humans migrated from Africa to other continents they carried these parasites wherever climate conditions allowed parasite transmission from host to host. Other parasites, however, were acquired throughout human biological and social evolutive history when new territories were occupied. Paleoparasitology data is a valuable source to recover emergence and disappearance of parasite infections through analysis of archaeological remains. Parasites can be used as biological markers of prehistoric human migrations. They are also indicators of diet, as parasite life cycles are related to specific kinds of food consumed by human groups in the different habitats they occupied. We review paleoparasitological findings in South America, comparing human-host and intestinal parasites with life conditions and environmental relationships through time.

Key words: Paleoparasitology, coprolites, mummies, infectious diseases, ancient diseases, parasite-human evolution.

Algunos parásitos humanos se originaron en ancestrales prehomínidos de Africa. Especies de nemátodos tales como Enterobius vermicularis y Trichuris trichiura son compartidos por los humanos y otros primates filogenéticamente emparentados (Pan and Gorilla), lo cual indica que tales parásitos antiguamente infectaron a un ancestro común de estos grupos. Cuando los humanos migraron desde Africa a otros continentes, llevaron consigo parásitos a lugares donde las condiciones climáticas permitieran su transmisión de huésped a huésped. Sin embargo, otros parásitos fueron adquiridos a través de la historia evolutiva humana biológica y social, durante la ocupación de nuevos territorios. Las evidencias paleoparasitológicas, encontradas a través del análisis de material arqueológico, son una fuente valiosa para recuperar información de infecciones parasitarias emergentes y desaparecidas. Los parásitos pueden ser usados como marcadores biológicos de migraciones humanas prehistóricas. Además de esto, son indicadores de dieta, debido a que los ciclos de vida parasitarios están relacionados con ciertos alimentos que fueron consumidos por grupos humanos en los diferentes hábitats ocupados. El objetivo de esta revisión bibliográfica es reunir los hallazgos paleoparasitológicos en Sudamérica y comparar la relación huésped humano y parásitos intestinales con condiciones de vida y relaciones ambientales a través del tiempo.

Palabras claves: paleoparasitología, coprolitos, momias, enfermedades infecciosas, enfermedades antiguas, evolución parásito-humano.

Parasites have been found in ancient remains throughout the world (Bouchet et al. 2003; Gonçalves et al. 2003). Coprolites and latrine contents are the main source of intestinal parasite

eggs. Protozoan cysts, however, are not as easy to identify. Nevertheless, serologic techniques have been successfully applied to identify *Giardia duodenalis*, *Giardia intestinalis* and *Entamoeba histolytica*

* Este artículo fue presentado en el Taller de Bioarqueología "Avances y Desafíos del Paleoambiente y la Paleoparasitología", Arica, noviembre 2008. De 15 ponencias expuestas 7 fueron seleccionadas para ser evaluadas por investigadores externos. Luego fueron editadas por Bernardo Arriaza y Nancy Orellana, en su calidad de editores invitados en este número.

¹ Escola Nacional de Saúde Pública, Fundação Oswaldo Cruz, Rua Leopoldo Bulhões 1480, CEP 21041-210, Rio de Janeiro, Brasil. adauto@ensp.fiocruz.br; dleles03@yahoo.com; lucianasianto@gmail.com

² University of Nebraska, Lincoln, Nebraska, Estados Unidos. kreinhard1@unl.edu

³ Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Av. Brasil 4365, CEP 21045-900, Rio de Janeiro, Brasil. alena@ioc.fiocruz.br

⁴ CONICET-Universidad Nacional de Mar del Plata, Argentina. mhufugassa@hotmail.com

⁵ Instituto de Alta Investigación, Universidad de Tarapacá, Arica, Chile. barriazaarica@gmail.com

⁶ Convenio de Desempeño-UTA/Mineduc, Universidad de Tarapacá, Arica, Chile.

⁷ Departamento de Antropología, Universidad de Tarapacá, Arica, Chile.

⁸ Universidad de la Cordillera, La Paz, Bolivia. nancy.orellana@gmail.com

⁹ Museo Nacional de Etnografía y Folklore, La Paz, Bolivia.

antigens in human coprolites (Gonçalves et al. 2002, 2004; Le Bailly et al. 2008). Other parasites, such as the protozoan *Trypanosoma cruzi* was recovered from mummified tissues and bone fragments by molecular biology techniques (Aufderheide et al. 2004; Fernandes et al. 2008; Ferreira et al. 2000; Guhl et al. 1999, 2000; Lima et al. 2008).

Parasitology is relevant to anthropology and the impact of parasites on human culture can be seen in language, art, religion, biology and archaeology (Araújo et al. 2003). Parasites and parasitic disease have cultural expressions as represented in Moche ceramic art (Heck 2004). The use of regular parasitological techniques, modified and adapted to paleoparasitological research increased and expanded parasite findings to other kinds of archaeological remains (Fugassa et al. 2006; Fugassa, Sardella et al. 2008; Harter et al. 2003; Harter-Lailheugue and Bouchet 2006).

History of Paleoparasites Found on Archaeological Remains

Since the first studies in South American human coprolites (Pizzi and Schenone 1954), paleoparasitology has developed as a new research line and this new science appeared based on solid data founded on parasite findings from well-dated and well-identified context, as predicted by Cockburn (1967). Common wisdom about human health conditions and parasite infections in the New World was challenged by paleoparasitological finds. Hookworm (ancylostomids), whipworm (*Trichuris trichiura*), pinworm (*Enterobius vermicularis*), and roundworm (*Ascaris lumbricoides*) were found infecting prehistoric populations both in North and South America.

Paleoparasitology focuses on the recovery of parasites from archaeological sites to further elucidate the ancient impacts of parasites on culture. Tracing the origins of parasites is one research thread of importance to paleopathology (Reinhard 1990). Parasitism has a central role in bioarchaeology (Reinhard 2008; Reinhard and Bryant 2008) and has long been recognized as a tool for the archaeologist (Reinhard 1992).

As reviewed by Reinhard and Bryant (2008), the relation between cultural development and the nature of parasitism has long been investigated. Reinhard (1988) first formalized this area of research in the Southwestern United States by testing Cockburn's

(1971) hypotheses about the impact of agriculture on parasitism. Reinhard showed that zoonotic infections, long standing in ancient populations, nearly disappeared in horticulturalists as human-specific parasites emerged. Martinson et al. (2003), Reinhard and Buikstra (2003) and Santoro et al. (2003) continued to explore the impact of parasitic disease ecology in the Andes. They clearly defined the role of trade, dress, and water contamination on parasitic disease among the Chiribaya of Peru (Martinson et al. 2003; Reinhard and Buikstra 2003). Santoro et al. (2003) demonstrated that empire development dramatically modified parasite ecology of the Lluta Valley, Chile, during Inka expansion to this lowland territory.

Reinhard (1992) noted that the most sensational contribution of paleoparasitology was towards a better understanding of the first human migrations to the New World. In addition, Araújo et al. (1981), Araújo, Reinhard et al. (2008), and Montenegro et al. (2006) stated that human coastal and transoceanic migrations also need consideration because helminth could not survive the colder Beringia climate. Further work by Reinhard and Bryant (2008) introduced the concept of pathoecology, which links parasitism to diet, environmental stress, and other culturally-defined behaviors in the definition of ancient disease.

Molecular biology techniques were also applied in diagnosing intestinal parasites. Loreille et al. (2001) and Loreille and Bouchet (2003) discussed the origin and evolution of *Ascaris lumbricoides* and *Ascaris suum* parasitism based on DNA studies performed on parasite eggs recovered from mediaeval latrines in Europe. Recently, Leles et al. (2008) applied molecular biology techniques and were able to recover ancient *Ascaris lumbricoides* DNA sequences from South American samples, negative by microscopy. These findings illustrate the importance of molecular biology techniques in diagnosing parasites in ancient remains. Molecular biology techniques allow the study of genetic evolution of parasites and the timing of their introduction into human populations (Dittmar et al. 2006; Dittmar 2009). By recovering aDNA (ancient DNA) sequences and genotyping parasites from human remains increases the possibility to reconstruct the early dispersion patterns of parasites (Raoult and Drancourt 2008).

This paper summarizes the extent of the knowledge of intestinal parasite remains from South

America. As such, it familiarizes the reader with the current findings of parasites from this ecologically diverse and culturally complex area. We review intestinal parasite infections in pre-Columbian South America. Paleoparasitological data is used by archaeologists and anthropologists to reconstruct ancient events based on parasite life cycles and biological requirements to maintain transmission from host to host. Climate changes are also associated with the presence of parasite infections in ancient people. For this purpose, paleoparasitological data gathered by the laboratories of Fundação Oswaldo Cruz, Universidad Nacional de Mar del Plata, and Universidad de Tarapacá-Arica, as well as data from other sources published in the literature were reviewed. The goal of this paper is to discuss paleoparasitology results in South America aiming to improve techniques and integrate data obtained by the laboratories involved in this matter.

Techniques Used to Analyse Bioarchaeological Material

Organic remains were examined by parasitological regular techniques after rehydration using a trisodium phosphate aqueous solution (Na_3PO_4 -0.5%). According to the nature of the sample, different techniques are recommended. For most coprolite analyses, spontaneous sedimentation in glass jars is recommended (Araújo et al. 1998; Lutz 1919). To prevent bacterial and fungal growth or taphonomic processes that may cause DNA degradation, thus interfering in molecular biology technique that may be applied, all the remaining sediment in the jars should be preserved refrigerated (Pruvost et al. 2007).

In some archaeological contexts, coprolites are the main source for paleoparasitology analysis. For example, in northeastern Brazil coprolites are found dispersed in archaeological layers under rock-shelters or inside caves (Bouchet et al. 2003). Mummified bodies are rarely found in Brazilian archaeological sites. According to Mendonça de Souza (pers. comm. 2008) preservation of the dead was never a common practice among Brazilian indigenous cultures because they believe the soul should be released soon after death. Just after death the individual was cremated, buried, or sometimes the ashes were eaten by others (Chagnon 1983). Paleoparasitological research in Brazil started with an interest in knowing what parasite infections existed

before the coming of Europeans and Africans to the Americas (Ferreira et al. 1988). These results were used to trace prehistoric migration routes for the peopling of the continent (Araújo et al. 1981; Araújo, Reinhard et al. 2008; Araújo, Reinhard and Ferreira 2008; Reinhard et al. 2001). Wild animal coprolites were also examined from many archaeological sites, especially focused on parasites of animals that may infect humans (Chame 2003; Sianto et al. 2005). Due to the collaboration with the University of Nebraska, Lincoln, ancient food remains turned to be also a research line at Fundação Oswaldo Cruz (FIOCRUZ). Molecular paleoparasitology has a special laboratory to develop techniques and diagnoses in conditions avoiding modern DNA contamination. The Laboratory of Paleoparasitology at FIOCRUZ, also has an image bank that will soon be available on the Internet.

A considerable amount of American paleoparasitological data appeared recently in the literature. Even Southern Patagonian coprolites were collected and examined (Fugassa 2007, 2008; Fugassa, Bayer and Sardella 2008). However as climatic conditions in Patagonia are not conducive to preservation, organic materials are continuously exposed to extreme dryness and cold and sometimes to water percolation. Due to the scarcity of coprolites and other organic material a microscopic method was developed to search for parasites at the Laboratory of Paleoparasitology of the Universidad Nacional de Mar del Plata, Argentina (Fugassa et al. 2006). The method used sediments recovered from the pelvic girdle and proved to be effective; it allowed for recovery of parasite eggs from human sacral foramina stored in museum and scientific institutions (Jones 1982; Fugassa, Denegri et al. 2006; Reinhard 1992). The main reason for pursuing paleoparasitological research in Argentina is to understand the impact of disease during the paleoepidemiological transition caused by the contact of Europeans with Patagonian Aborigines.

At the Universidad Nacional Mayor de San Marcos in Lima, Perú, Inés Gárate and her team applied regular parasitological techniques to examine human coprolites excavated in the archaeological site of Caral to search for parasites (Gárate et al. 2005). Caral is the oldest city of the western hemisphere, dated of 5000 years before present (BP) (Solis et al. 2001). Paleoparasitological research started recently exploring this archaeological site material, and first results are expected to be published soon. Paleoparasitology in Peruvian archaeological remains

has a tradition in coprolite (Patrucco et al. 1983) and mummy analysis (Fornaciari et al. 1992; Salo et al. 1994), but also on ectoparasites (Rick et al. 2002; Dittmar 2000; Dittmar et al. 2003; Raoult et al. 2008). Another group interested in paleoparasitological research was recently created by Luis Huamán and Hugo Flores, and their students at the Universidad Cayetano Heredia. The main interest of these specialists is plant and food remains and parasite associated with food resources and diet.

Regarding the study of parasites in mummified bodies, the Chinchorro mummies of Northern Chile present an interesting challenge. The most ancient mummies were prepared by defleshing the body to the bones and taking out abdominal viscera so intestinal contents are not expected to be found. Desert climate is especially suitable to preserve coprolites and paleoparasites. Fouant et al. (1982) reported a low incidence of parasites in Chilean and Peruvian mummies. Considering the tradition of paleoparasitological research in Chile, the interests varied from tuberculosis and Chagas disease (Rothhammer et al. 1985), intestinal parasites, and diet as well as implications of ENSO phenomenon (Arriaza et al. 2010; Williams et al. 2008).

In all the laboratories involved in paleoparasitological research, ectoparasites have been studied. Mummies of humans and other animals have been searched for lice, fleas, and other arthropods. It is important to look for parasites in the wrapping tissues of the mummies and collect sediment samples near the body.

Intestinal Parasites

Paleoparasitological research proved that common intestinal parasites infected pre-Columbian people long before European and African migrations (Gonçalves et al. 2003). Data also showed that Amerindians acquired infections by eating intermediate hosts or being in close contact with vectors and hosts of parasites that infect humans (Patrucco et al. 1983; Sianto et al. 2005).

Some intestinal parasites that originated in hominid African ancestors were introduced into South America with their human hosts long before European conquest (Araújo, Reinhard, Ferreira et al. 2008). Prehistoric people who lived on the American continent were infected with common bacteria, protozoa, and helminths as in other parts of the world. Although hookworms, pinworm,

whipworm, and roundworm infection were the same, prevalence rates among pre-Columbian populations and transmission dynamics should behave in a different way according to the different cultures where they were infecting. Parasite transmission dynamics was not the same in the city of Caral, in Peru, or in the Atacama Desert where the Chinchorros lived, and in the semi-arid region of Brazilian northeast where hunter-gathering strategy was used by pre-historic groups. Thus, generalization on parasites and ecology may lead to erroneous interpretations (Figure 1 and Table 1). Brazilian groups were numerous and lived in crowded villages before the European continent conquest (Heckenberger et al. 2008). Contrary to what is the common knowledge, Amerindian groups living in the Brazilian coast where numerous and just before European conquest some groups were displaced to the interior after tribal wars, and in turn exterminated small groups of hunter-gatherers living in caves and rock-shelters (Medeiros 2002).

Studies about the impact of European contact with indigenous population opened a new line of research, especially regarding paleoepidemiological transition unleashed by European arrival (Guichón et al. 2006). Fugassa, Cicchino et al. (2008) showed that hunter-gatherer groups of Patagonia would have had a parasitological profile, zoonosis caused by close wild animals contact or ingestion of intermediate hosts meat and vegetables contaminated with parasite eggs or cysts, as well as arthropods, such as oribatids, which are intermediate hosts for anoplocephalids. After European contact parasite infections and relative abundance of the common human intestinal worms *Ascaris lumbricoides* and *Trichuris trichiura* changed significantly (Fugassa et al. 2006). Crowding indigenous populations in religious missions, reducing mobility of the nomadic groups, and the fast and abundant arrival of domestic animals, mainly sheeps, caused a rapid change of the helminth and protozoan fauna, due to human/animal parasite cross infections. European also caused a reduction of local fauna. Europeans also introduced food storage, attracting rodents and their parasites to dwellings (Guichón et al. 2006).

Trichuris trichiura (whipworm) infection was common in South America on both sides of the Andean cordillera (Gonçalves et al. 2003). Whipworm eggs were the first to be found in a South American mummy, together with *Entamoeba coli* cysts (Pizzi and Schenone 1954). *Enterobius*




Figure 1. Map of South America showing the location of paleoparasitological findings listed in Table 1.
 Mapa de Sudamérica, muestra la localización de los paleoparásitos hallados, citados en la Tabla 1.

Table 1. Summary of South American paleoparasitological findings.
Resumen de los hallazgos paleoparasitológicos en Sudamérica.

Country	South America				
	Archaeological Site	No	Date	Parasites	Reference
Argentina	Pie de Palo	1	not available	<i>Enterobius vermicularis</i>	Fugassa et al. 2006
	Perito Moreno, Santa Cruz	2	6,540±110 BP	<i>Enterobius vermicularis</i> <i>Rabditoid larvae</i> <i>Trichuris</i> sp.	Fugassa 2008
	Nombre de Jesús, Cabo Virgenes	3	not available	<i>Ascaris lumbricoides</i>	Fugassa et al. 2006
	Orejas de Burro, Santa Cruz	4	3,720-3,978 BP	<i>Ascaridid</i>	Fugassa 2006
	Valle Encantado, Neuquén	5	1,000-500 BP	<i>Ancylostomid</i>	Gonçalves et al. 2003
	Las Mandibulas, Tierra del Fuego	6	Historic	<i>Capillarids</i>	Fugassa et al. 2008
	Caleta Falsa, Tierra del Fuego	7	850 years BP	<i>Capillarids</i>	Fugassa et al. 2008
	Cerro Norte XI, Pali Aike, Santa Cruz	8	not available	<i>Capillarids</i>	new findings
	Salitroso, Santa Cruz	9	not available	<i>Trichuris trichiura</i>	new findings
	Parador Nativo, Rio Negro	10	1,513±48 BP	<i>Trichuris trichiura</i>	Fugassa et al. 2006
	Centro Minero, Rio Negro	11	689±44 BP	<i>Trichuris trichiura</i>	Fugassa et al. 2006
Peru	Huamey Valley	12	1,000 AD	<i>Trichuris trichiura</i>	Patrucco et al. 1983
	Coast (southern)	13	10,000-4,000 BP	<i>D. pacificum</i>	Patrucco et al. 1983
	Osmore, Peru	14	1,020-1,476 BP	<i>D. pacificum</i>	Patrucco et al. 1983
	Huaca Prieta	15	3,000 BP	<i>Diphyllobotrium</i> sp.	Callen and Cameron 1960
Brazil	Toca do Meio, Piauí	16	8,800±60 BP	<i>Ascaris lumbricoides</i>	Leles et al. 2008
	Lapa Pequena, Minas Gerais	17	8,000-7,000 BP	<i>Ascaris lumbricoides</i>	Gonçalves et al. 2003
	Santa Elina, Mato Grosso	18	4,000-2,000 BP	<i>Ascaris lumbricoides</i> <i>Hymenolepis nana</i>	Gonçalves et al. 2003
	Gruta do Gentio II, Minas Gerais	19	3,490±120-4,30±70 BP	<i>Ascaris lumbricoides</i>	Leles et al. 2008
	Gentio Cave, Minas Gerais	20	3,490±120-430±70 BP	<i>Ascaris lumbricoides</i> <i>Acantocephala</i>	Gonçalves et al. 2003
	Pedra Furada, Piauí I	21	not available	<i>Trichuris trichiura</i>	Gonçalves et al. 2003
	Boqueirão Soberbo, Minas Gerais	22	4,905±85-1,325±60 BP	<i>Trichuris trichiura</i> <i>Acantocephala</i>	Ferreira et al. 1984
	Estrago Cave, Pernambuco	23	2,000 BP	<i>Trichuris trichiura</i>	Gonçalves et al. 2003
	Itacambira, Minas Gerais	24	not available	<i>Trichuris trichiura</i>	Confalonieri 1988
	Sítio do Meio, Piauí	25	not available	<i>Ancylostomid</i>	Gonçalves et al. 2003
Chile	Tiliviche, Iquique	26	4,110-1,950 BC	<i>D. pacificum</i> <i>Enterobius vermicularis</i> <i>Ancylostomid</i>	Ferreira et al. 1984 Gonçalves et al. 2003
	Tulán, San Pedro de Atacama	27	1,000 BC	<i>Trichuris trichiura</i>	Ferreira et al. 1984
	Lluta Valley, Arica	28	1,200-1,500 AD (Inka period)	<i>Trichuris trichiura</i> <i>Hymenolepis nana</i>	Santoro et al. 2003
	Toconao, San Pedro de Atacama	29	2,500-2,100 BP	<i>Ancylostomid</i>	Gonçalves et al. 2003
	El Plomo, Santiago	30	Pre-Columbian	<i>Trichuris trichiura</i>	Ferreira et al. 1984
	Catarpe, San Pedro de Atacama	31	1,450-1,525 AD	<i>Trichonstrongylus</i> sp.	Gonçalves et al. 2003
	San Miguel de Azapa	32	4,000-5,000 AP	<i>Diphyllobotrium</i> sp.	Reinhard and Urban 2003

vermicularis (pinworm) eggs, curiously, were only found among ancient populations who lived on the Pacific side. Molecular biology studies, however, showed interesting results, with two different pinworm lineages in the Pacific coast populations. One had the same sequences of North American lineages, but the other showed divergent nucleotides in some sequences (Iñiguez et al. 2003). According to the authors, this may be interpreted as the consequence of different origins. While one was introduced by human pre-historic migrations that crossed the Bering Land Bridge more than 20,000 years ago, the other lineage suggests an introduction by another route, in this case supporting transpacific contacts of Asian people in pre-Columbian times (Araújo, Reinhard and Ferreira 2008; Iñiguez et al. 2006).

Regarding intestinal protozoan infection, South American paleoparasitology data is not as representative, although the two most prevalent protozoan parasites found in coprolites and sediments were *Entamoeba histolytica* and *Giardia intestinalis* (Gonçalves, Araújo et al. 2002; Gonçalves, da Silva et al. 2004). Antigens of *Giardia* sp., *Cryptosporidium* sp., *Cyclospora* sp., and *Helicobacter pylori*, were identified in Andean mummies dated up to 3,000 years by fluorescent microscopy (Allison et al. 1999; Ortega and Bonavia 2003).

As evidenced by paleoparasitology finds, common human intestinal parasites were infecting South American pre-Columbian populations. Although data are still missing for many South American countries, an interesting paleoparasitological picture has been traced from Argentinean, Bolivian, Brazilian, Chilean, and Peruvian archaeological material (Ferreira et al. 2008). When these findings are compared with North American paleoparasitology, a similar picture appears and human strategies to confront different environments can be seen between the continents (Reinhard 1992).

Many of these parasites originated in Old World human populations (Araújo and Ferreira 1995). They came out of Africa with the peopling of Europe. From there, they spread through the world where climate conditions allowed transmission (Araújo, Reinhard and Ferreira 2008). However, humans acquired other parasites from animals whenever humans invaded new habitats, adopted new habits, or created new technologies amplifying their range of influence on the environment.

Humans adopted food habits according to the occupied region. This is evidenced by coprolite

paleoparasitological analysis. *Diphyllobothrium pacificum* was a common intestinal infection in the past as it is today in the Pacific coast. The consumption of raw fish is a common habit in many countries and different cultures all over the world. Baer et al. (1967) reported the presence of *D. pacificum* in modern Peruvian coast population, later *D. pacificum* infection was confirmed in Peruvian and Chilean pre-Columbian coprolites (Araújo et al. 1983; Ferreira et al. 1984; Patrucco et al. 1983; Reinhard and Urban 2003). *Diphyllobothrium pacificum* human transmission is due to the consumption of raw fish containing parasites.

Echinostoma sp. eggs were found in a Brazilian mummy (Sianto et al. 2005). As this parasite was not recorded in modern South American population, the finding was carefully investigated before publishing this information. It is interesting to note that this case of *Echinostoma* human infection in pre-historic South America was associated with a case of Chagas disease, with intestinal lesions characterized by megacolon (Fernandes et al. 2008). In this case two important points should be emphasized: firstly, the description of Chagas disease lesions in a mummy outside the Andean region, confirmed by molecular paleoparasitology studies (Fernandes et al. 2008); secondly, an infection by *Echinostoma* in South American human population not yet described (Sianto et al. 2005). These findings point to the need of further detailed studies at this archaeological site.

Animal Parasites in Humans

The presence of animal parasites in human coprolites shows the use of different food resources. However, these habits may still be maintained today by traditional groups. For example, Coimbra Jr. and Mello (1981) found *Capillaria* eggs in modern Amazonian Indian group, and Fugassa, Bayer and Sardella (2008) identified *Capillaria* eggs in animal coprolites in Patagonia, which may point to a potential source of infection for humans in this region. *Capillaria* species rarely infect humans, but eggs can pass with feces when humans eat an infected host. Thus, the eggs can be found in human feces, but disappear after a few days.

Fugassa et al. (2006) presented an overview of human and animal intestinal helminths and protozoa found in archaeological material excavated in Patagonia. These findings illustrate the diversity

of parasites that can be found in archaeological remains, in close contact with humans beginning 10,000 years ago. Therefore, paleoparasitology is a powerful tool to investigate the variety of food sources used by humans at different times by studying organic remains preserved in archaeological sites.

Ectoparasites

South American mummies provided positive results for ectoparasites. In the fur of mummified guinea pigs (*Cavia aperea* f. *porcellus*), numerous well-preserved ectoparasites (lice, fleas, mites) could be recovered (Dittmar 2002; Dittmar et al. 2003). Moreover, paleoparasitological examination of Amerindian clothes provided parasite finds, such as ectoparasites (Fugassa et al. 2007). Molecular biology studies were also performed (Dittmar et al. 2003; Raoult et al. 2008), as well as paleoepidemiological and quantitative attempts to evaluate head lice infection (Reinhard and Buikstra 2003). Head lice were found still attached to hair fragments of a skeleton dated to 10,000 years in northeastern Brazilian (Araújo et al. 2000). Crablouse was also found infecting Peruvian mummies (Rick et al. 2002). Rivera et al. (2008) reviewed the findings of head lice in

South American mummies recording new finds of *Anthropophthirus capitis* (a new genus proposed by Retana-Salazar and Ramirez-Morales 2006 to substitute *Pediculus humanus capitis*) in mummies of the Chinchorro tradition dated to 4,000 years. This new species proposal is very controversial, but it was proposed due to genetic similarities found between *Pediculus humanus* (human body lice) and the *Parapediculus* (ectoparasite of South American monkeys).

To conclude, South American paleoparasitologists, together with North American colleagues, are building an evolving picture of parasite infections dated from the peopling of the Americas (Araújo, Reinhard and Ferreira 2008; Araújo, Reinhard et al. 2008). Paleoparasitological data, through the use of microscopic, immunology and molecular techniques, is an efficient way to provide information about human behavior in the past, contributing to understanding the origin and evolution of infectious diseases and their impact on prehistoric populations.

Acknowledgements: Supported by the Brazilian agencies CNPq, CAPES, and FAPERJ. We also want to thank the Convenio Desempenho UTA-MINEDUC for their support and the manuscript editors, and outside reviewers for their comments and suggestions.

References Cited

- Allison, M.J., T. Bergman, and E. Gerszten
1999 Further studies on fecal parasites in antiquity. *American Journal of Clinical Pathology* 112(5):605-609.
- Araújo, A., and L.F. Ferreira
1995 Oxiurias e migrações pré-históricas. *Historia, Ciências, Saude - Manguinhos* 2(1):99-109.
- Araújo, A., L.F. Ferreira, and U. Confalonieri
1981 A contribution to the study of helminth findings in archaeological material in Brazil. *Revista Brasileira de Biologia* 41:873-881.
- Araújo, A., L.F. Ferreira, U.E. Confalonieri, L. Nuñez, and F.O. Cruz
1983 Eggs of *Diphyllobothrium pacificum* in precolumbian human coprolites. *Paleopathology Newsletter* 41:11-13.
- Araújo, A., L.F. Ferreira, N. Guidon, N. Maues Da Serra Freire, K.J. Reinhard, and K. Dittmar
2000 Ten thousand years of head lice infection. *Parasitology Today* 16(7):269.
- Araújo, A., A.M. Jansen, F. Bouchet, K. Reinhard, and L.F. Ferreira
2003 Parasitism, the diversity of life, and paleoparasitology. *Memórias do Instituto Oswaldo Cruz* 98:5-11.
- Araújo, A., K. Reinhard, O.M. Bastos, L.C. Costa, C. Pirmez, A. Iñiguez, A.C. Vicente, C.M. Morel, and L.F. Ferreira
1998 Paleoparasitology: perspectives with new techniques. *Revista do Instituto de Medicina Tropical de Sao Paulo* 40(6):371-376.
- Araújo, A., K. Reinhard, and L.F. Ferreira
2008 Parasite findings in archaeological remains: diagnosis and interpretation. *Quaternary International* 174:1-4, 180:17-21.
- Araújo, A., K.J. Reinhard, L.F. Ferreira, and S.L. Gardner
2008 Parasites as probes for prehistoric human migrations? *Trends in Parasitology* 24:112-115.
- Arriaza, B., K. Reinhard, A. Araújo, N. Orellana, and V. Standen
2010 Possible influence of the ENSO phenomenon on the pathoecology of diphyllobothriasis and anisakiasis in ancient Chinchorro populations. *Memórias do Instituto Oswaldo Cruz* 105(1):66-72.
- Aufderheide A.C., W. Salo, M. Madden, J. Streit, J. Buikstra, F. Guhl, B. Arriaza, C. Renier, L.E. Wittmers Jr., G. Fornaciari, and M. Allison
2004 A 9,000-year record of Chagas' disease. *Proceedings of the National Academy of Sciences* 101(7):2034-2039.

- Baer, J.G., H. Miranda, W. Fernandez, and J. Medina
1967 Human diphyllbothriasis in Peru. *Zeitschrift für Parasitenkunde* 28(3):277-289.
- Bouchet F, N. Guidon, K. Dittmar, S. Harter, L.F. Ferreira, S.M. Chaves, K. Reinhard, and A. Araújo
2003 Parasite remains in archaeological sites. *Memorias do Instituto Oswaldo Cruz* 98(1):47-52.
- Callen, O., and T.W.M. Cameron
1960 A prehistoric diet as revealed in coprolites. *The New Scientist* 8:35-40.
- Chagnon, N.A.
1983 *Yanomamo: The Fierce People*. CBS College Publishing, New York, NY.
- Chame, M.
2003 Terrestrial mammal feces: A morphometric summary and description. *Memórias do Instituto Oswaldo Cruz* 98(Suppl. I):71-94.
- Cockburn, A.
1967 *Infectious Diseases: Their Evolution and Eradication*. Charles C. Thomas, Springfield.
1971 Infectious diseases in ancient populations. *Current Anthropology* 12:45-62.
- Coimbra Jr., C.E.A., and D.A. Mello
1981 Enteroparasitas e *Capillaria sp.* entre o grupo Suruí, Parque Indígena Aripuanã, Rondônia. *Memórias de Instituto Oswaldo Cruz* 76:299-302.
- Confalonieri, U.
1988 Paleoepidemiologia de *Trichuris trichiura* na América. In *Paleoparasitologia no Brasil*, edited by L.F. Ferreira, A. Araújo, and U. Confalonieri, pp. 120-137. Editora PEC/ENSP, Rio de Janeiro.
- Dittmar, K.
2000 Evaluation of ectoparasites on the Guinea pig mummies of El Yaral and Moquegua Valley, in Southern Peru. *Chungara Revista de Antropología Chilena* 32:123-125.
2002 Arthropod and helminth parasites of the wild guinea pig, *Cavia aperea*, from the Andes and the Cordillera in Peru, South America. *Journal of Parasitology* 88: 409-411.
2009 Old parasites for a New World: The future of paleoparasitological research. A review. *Journal of Parasitology* 95:365-371.
- Dittmar, K., U. Mamat, M. Whiting, T. Goldmann, K. Reinhard, and S. Guillen
2003 Techniques of DNA-studies on prehispanic ectoparasites (*Pulex sp.*, Pulicidae, Siphonaptera) from animal mummies of the Chiribaya culture, Southern Peru. *Memórias do Instituto Oswaldo Cruz* 98:53-58.
- Dittmar, K., S.M. de Souza, and A. Araújo
2006 Challenges of phylogenetic analyses of aDNA sequences. *Memorias do Instituto Oswaldo Cruz* 101:9-13.
- Fernandes, A., A.M. Iñiguez, V.S. Lima, S.M. de Souza, L.F. Ferreira, A.C. Vicente, and A.M. Jansen
2008 Pre-Columbian chagas disease in Brazil: *Trypanosoma cruzi* I in the archaeological remains of a human in Peruaçu Valley, Minas Gerais, Brazil. *Memórias do Instituto Oswaldo Cruz* 103:514-516.
- Ferreira L.F., A. Araújo, and U. Confalonieri
1988 *Paleoparasitologia no Brasil*. Ed. PEC/ENSP-FIOCRUZ, Rio de Janeiro.
- Ferreira, L.F., A.J.G. Araújo, and U.E.C. Confalonieri
1982 The finding of helminth eggs in a Brazilian mummy. *Tropical Medicine & Hygiene* 77(1):65-67.
- Ferreira, L.F., A.J. de Araújo, U.E. Confalonieri, and L. Nuñez
1984 The finding of eggs of *Diphyllbothrium* in human coprolites (4,100-1,950 B.C.) from Northern Chile. *Memórias do Instituto Oswaldo Cruz* 79:175-180.
- Ferreira, L.F., C. Britto, M.A. Cardoso, O. Fernandes, K. Reinhard, and A. Araújo
2000 Paleoparasitology of Chagas disease revealed by infected tissues from Chilean mummies. *Acta Tropica* 75:79-84.
- Ferreira, L.F., K. J. Reinhard, and A. Araújo
2008 *Paleoparasitologia*. Ed. Fiocruz, Rio de Janeiro, Brasil.
- Fornaciari, G., M. Castagna, P. Viacava, A. Tognetti, G. Bevilaqua, and E.L. Segura
1992 Chagas' disease in Peruvian Inca mummy. *Lancet* 339:128-129.
- Fugassa, M.H.
2006 *Enteroparasitosis en Poblaciones Cazadoras-Recolectoras de Patagonia Austral*. Tesis Doctoral. Universidad Nacional de Mar del Plata, Buenos Aires.
2007 Camélidos, parásitos y ocupaciones humanas: registros paleoparasitológicos en Cerro Casa de Piedra 7 (Parque Nacional Perito Moreno, Santa Cruz, Argentina). *Intersecciones en Antropología* 8:265-269.
2008 Aportes paleoparasitológicos para el sitio arqueológico CCP7, P. N. Perito Moreno, Santa Cruz. *Cuadernos del Instituto Nacional de Antropología y Pensamiento Latinoamericano* 21:263-265.
- Fugassa, M.H., A. Araújo, and R.A. Guichón
2006 Quantitative paleoparasitology applied to archaeological sediments. *Memórias do Instituto Oswaldo Cruz* 101:29-33.
- Fugassa, M.H., M.S. Bayer, and N.H. Sardella
2008 Examen paleoparasitológico de coprolitos de felinos en Patagonia (Holoceno Medio). Manuscript in possession of the authors.
- Fugassa M. H., A. Cicchino, N. H. Sardella, R. A. Guichón, G. M. Denegri, and A. Araújo
2007 Nuevas fuentes de evidencia para la paleoparasitología y la antropología biológica en Patagonia. *Revista de la Asociación de Antropología Biológica* 9:51-57.
- Fugassa, M.H., A., Cicchino, N. H. Sardella, R.A. Guichón, G.M. Denegri, and A. Araújo
2008 Nuevas fuentes de evidencia para la paleoparasitología y la antropología biológica en Patagonia. *Revista Argentina de Antropología Biológica* 9:51-57.
- Fugassa, M.H., G.M. Denegri, N.H. Sardella, A. Araújo, R.A. Guichón, P.A. Martinez, M.T. Civalero, and C. Aschero
2006 Paleoparasitological records in canid coprolite from Patagonia, Argentina. *Journal of Parasitology* 92:1110-1113.
- Fugassa, M.H., N.H. Sardella, R.A. Guichón, G.M. Denegri, and A. Araújo
2008 Paleoparasitological analysis applied to museum-curated sacra from meridional Patagonian collections. *Journal of Archaeological Science* 35:1408-1411.

- Fugassa, M.H., V. Taglioretti, M.L.C. Gonçalves, A. Araújo, N.H. Sardella, and G.M. Denegri
2008 *Capillaria* spp. findings in Patagonian archaeological sites: Statistical analysis of morphometric data. *Memórias do Instituto Oswaldo Cruz* 103:104-105.
- Gárate, I., B. Suyo, M. Delgado, H. Solís, and P. Castellanos
2005 Hallazgo de nematode y huevos de *Ascaris* sp. en coprolitos de la momia "Shamana Alada", Cerrillos, Ica, Peru. *Boletín Chileno de Parasitología y de Parasitología al Día* 60 (T2):325.
- Gonçalves, M.L., A. Araújo, R. Duarte, J.P. da Silva, K. Reinhard, F. Bouchet, and L.F. Ferreira
2002 Detection of *Giardia duodenalis* antigen in coprolites using a commercially available enzyme-linked immunosorbent assay. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 96:640-643.
- Gonçalves, M.L., A. Araújo, and L.F. Ferreira
2003 Human intestinal parasites in the past: New findings and a review. *Memórias do Instituto Oswaldo Cruz* 98:103-118.
- Gonçalves, M.L., V.L. da Silva, C.M. de Andrade, K. Reinhard, G.C. da Rocha, M. Le Bailly, F. Bouchet, L.F. Ferreira, and A. Araújo
2004 Amoebiasis distribution in the past: First steps using an immunoassay technique. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 98:88-91.
- Guhl, F., C. Jaramillo, G.A. Vallejo, F. Cárdenas-Arroyo, and A. Aufderheide
2000 Chagas disease and human migration. *Memórias do Instituto Oswaldo Cruz* 95:553-555.
- Guhl, F., C. Jaramillo, G.A. Vallejo, R. Yockteng, F. Cárdenas-Arroyo, G. Fornaciari, B. Arriaza, and A.C. Aufderheide
1999 Isolation of *Trypanosoma cruzi* DNA in 4,000-year-old mummified human tissue from Northern Chile. *American Journal of Physical Anthropology* 108:401-407.
- Guichón, R.A., J.A. Suby, R. Casali, and M.H. Fugassa
2006 Health at the time of Native-European contact in Southern Patagonia: First steps, results, and prospects. *Memórias do Instituto Oswaldo Cruz* 101:97-105.
- Harter, S., M. Le Bailly, F. Janot, and F. Bouchet
2003 First paleoparasitological study of an embalming rejects jar found in Saqqara, Egypt. *Memórias do Instituto Oswaldo Cruz* 98:119-121.
- Harter-Lailheugue S., and F. Bouchet
2006 Etude paléoparasitologique d'éléments atypiques de la Basse et Haute Vallée du Nil. *Bulletin de la Société de Pathologie Exotique* 99:53-57.
- Heck, J.
2004 Krankheit und körperdeformation in darstellungen auf Moche-Tongefäßen analyse und synopse aus ärztlicher Sicht. *Baessler-Archiv* 52:105-124.
- Heckenberger, M.J., J.C. Russell, C. Fausto, J.R. Toney, M.J. Schmidt, E. Pereira, B. Franchetto, and A. Kuikuro
2008 Pre-Columbian urbanism, anthropogenic landscapes, and the future of the Amazon. *Science* 321:1214-1217.
- Iñiguez, A.M., K.J. Reinhard, A. Araújo, L.F. Ferreira, and A.C. Vicente
2003 *Enterobius vermicularis*: ancient DNA from North and South American human coprolites. *Memórias do Instituto Oswaldo Cruz* 98:67-69.
- Iñiguez, A.M., K. Reinhard, M.L. Carvalho Gonçalves, L.F. Ferreira, A. Araújo, and A.C. Paulo Vicente
2006 SL1 RNA gene recovery from *Enterobius vermicularis* ancient DNA in pre-Columbian human coprolites. *International Journal for Parasitology* 36:1419-1425.
- Jones, A.
1982 Recent finds of intestinal parasites ova at York, England. Papers on Paleopathology. 4th European Members Meeting, Middelburg, Antwerpen, p. 7.
- Leles, D., A. Araújo, L.F. Ferreira, A.C. Vicente, and A.M. Iñiguez
2008 Molecular paleoparasitological diagnosis of *Ascaris* sp. from coprolites: new scenery of ascariasis in pre-Colombian South America times. *Memórias do Instituto Oswaldo Cruz* 103:106-108.
- Le Bailly, M., M.L. Gonçalves, S. Harter-Lailheugue, F. Prodéo, A. Araújo, and F. Bouchet
2008 New finding of *Giardia intestinalis* (Eukaryote, Metamonad) in Old World archaeological site using immunofluorescence and enzyme-linked immunosorbent assays. *Memórias do Instituto Oswaldo Cruz* 103:298-300.
- Lima, V., A. Iñiguez, K. Otsuki, L.F. Ferreira, A. Araújo, A. Vicente, and A. Jansen
2008 Chagas disease by *Trypanosoma cruzi* lineage I in hunter-gatherer ancient population in Brazil. *Emerging Infectious Diseases* 14:1001-1002.
- Loreille, O., and F. Bouchet
2003 Evolution of ascariasis in humans and pigs: a multi-disciplinary approach. *Memórias do Instituto Oswaldo Cruz* 98:39-46.
- Loreille, O., E. Roumat, O. Verneau, F. Bouchet, and C. Hänni
2001 Ancient DNA from *Ascaris*: extraction amplification and sequences from eggs collected in coprolites. *International Journal of Parasitology* 31:1101-1106.
- Lutz, A.
1919 O *Schistosomum mansoni* e a schistosomatose, segundo observações feitas no Brasil. *Memórias do Instituto Oswaldo Cruz* 11:121-150.
- Martinson, E., K.J. Reinhard, J.E. Buikstra, and K. Dittmar
2003 Pathoecology of Chiribaya parasitism. *Memórias do Instituto Oswaldo Cruz* 98:195-205.
- Medeiros, R.P.
2002 Povos indígenas do sertão nordestino no período colonial: descobrimento, alianças, resistência e encobrimento. *FUMDHamentos* 2:7-52.
- Montenegro, A., A. Araújo, M. Eby, L.F. Ferreira, R. Hetherington, and A. Weaver
2006 Parasites, paleoclimate and the peopling of the Americas: using the hookworm to time the Clovis migration. *Current Anthropology* 47:193-198.
- Ortega, Y.R., and D. Bonavia
2003 *Cryptosporidium*, *Giardia*, and *Cyclospora* in ancient Peruvians. *Journal of Parasitology* 89:635-636.
- Patrucco, R., R. Tello, and D. Bonavia
1983 Parasitological studies of coprolites of pre-hispanic Peruvian populations. *Current Anthropology* 24:393-394.
- Pizzi, T., and H. Schenone
1954 Hallazgo de huevos de *Trichuris trichiura* en el contenido intestinal de un cuerpo arqueológico arcaico Inca. *Boletín Chileno de Parasitología* 9:73-75.

- Pruvost, M., R. Schwarz, V.B. Correia, S. Champlot, S. Braguier, N. Morel, Y. Fernandez-Jalvo, T. Grange, and E.M. Geigl
2007 Freshly excavated fossil bones are best for amplification of ancient DNA. *Proceedings of the National Academy of Sciences* 104:739-744.
- Raoult, D., and M. Drancourt
2008 *Paleomicrobiology: Past Human Infections*. Springer-Verlag, Berlin Heidelberg.
- Raoult, D., D.L. Reed, K. Dittmar, J.J. Kirchman, J.M. Rolain, S. Guillen, and J.E. Light
2008 Molecular identification of lice from pre-Columbian mummies. *Journal of Infectious Diseases* 197:535-543.
- Reinhard, K.J.
1988 Cultural ecology of prehistoric parasitism on the Colorado Plateau as evidenced by coprology. *American Journal of Physical Anthropology* 77:355-366.
1990 Archaeoparasitology in North America. *American Journal of Physical Anthropology* 82:145-163.
1992 Parasitology as an interpretive tool in archaeology. *American Antiquity* 57:231-245.
2008 Parasite pathoecology of chacoan great houses: the healthiest and wormiest ancestral pueblos. In *Chaco's Northern Prodigies Salmon, Aztec, and the Ascendancy of the Middle San Juan Region after AD 1100*, edited by P.F. Reed, pp. 86-95. University of Utah Press, Salt Lake City.
- Reinhard, K., A. Araújo, L.F. Ferreira, and C.E. Coimbra
2001 American hookworm antiquity. *Medical Anthropology* 20:96-101.
- Reinhard, K.J., and V.M. Bryant
2008 Pathoecology and the future of coprolite studies in bioarchaeology. In *Reanalysis and Reinterpretation in Southwestern Bioarchaeology*, edited by A.W.M. Stodder, pp. 205-224. Arizona State University Press, Tempe.
- Reinhard, K., and J. Buikstra
2003 Louse infestation of the Chiribaya Culture, Southern Peru: variation in prevalence by age and sex. *Memórias do Instituto Oswaldo Cruz* 98:173-179.
- Reinhard, K., and O. Urban
2003 Diagnosing ancient diphyllbothriasis from Chinchorro mummies. *Memórias do Instituto Oswaldo Cruz* 98:191-193.
- Retana-Salazar, A.P., and R. Ramirez-Morales
2006 Establecimiento de un nuevo género de piojos (Phthiraptera: Pediculidae) asociado al hombre (Primates: Hominidae). *Brenesia* 65:61-70.
- Rick, F.M., G.C. Rocha, K. Dittmar, C.E. Coimbra Jr, K. Reinhard, F. Bouchet, L.F. Ferreira, and A. Araújo
2002 Crab louse infestation in pre-Columbian America. *Journal of Parasitology* 88:1266-1267.
- Rivera, M.A., K.Y. Mumcuoglu, R.T. Matheny, and D.G. Matheny
2008 Huevecillos de *Anthropophthirus capitis* en momias de la tradición Chinchorro, Camarones 15-D, Norte de Chile. *Chungara Revista de Antropología Chilena* 40:31-39.
- Rothhammer, F., M.J. Allison, L. Núñez, V. Standen, and B. Arriaza
1985 Chagas' disease in pre-Columbian South America. *American Journal of Physical Anthropology* 68:495-498.
- Salo, W., A. Aufderheide, J. Buikstra, and T. Holcomb
1994 Identification of Mycobacterium tuberculosis DNA in a pre-Columbian Peruvian mummy. *Proceedings of the National Academy of Sciences* 91:2091-2094.
- Santoro, C., S.D. Vinton, and K. Reinhard
2003 Inca expansion and parasitism in the Lluta Valley: preliminary data. *Memórias do Instituto do Oswaldo Cruz* 98:161-163.
- Sianto, L., K.J. Reinhard, M. Chame, S. Chaves, S. Mendonça, M.L. Gonçalves, A. Fernandes, L.F. Ferreira, and A. Araújo
2005 The finding of Echinostoma (Trematoda: Digenea) and hookworm eggs in coprolites collected from a Brazilian mummified body dated 600-1,200 years before present. *Journal of Parasitology* 91:972-975.
- Solis, R.S., J. Haas, and W. Creamer
2001 Dating Caral, a preceramic site in the Supe Valley on the central coast of Peru. *Science* 292:723-726.
- Williams A., C.M. Santoro, M.A. Smith, and C. Latorre
2008 The impact of ENSO in the Atacama Desert and Australian arid zone: exploratory time-series analysis of archaeological records. *Chungara Revista de Antropología Chilena* 40:245-259.