

Dynamics of soil organic carbon and dissolved organic carbon in *Robinia pseudoacacia* forests

H. Lv^{1,3} and Z. Liang^{1,2*}

¹Institute of Soil and Water Conservation, Chinese Academy of Science and Ministry of Water Resources, Yangling, Shaanxi 712100, P.R. China. ²College of Life Science, Northwest A&F University, Yangling, Shaanxi 712100, P.R. China. ³Weinan Teachers University, Weinan, Shaanxi 714000, P.R. China. *Corresponding author: liangzs@ms.iswc.ac.cn

Abstract

We investigated the variation patterns of organic carbon in soil and soil solution of four selected *Robinia pseudoacacia* forests aged 10a, 25a, 31a, and 35a, as well as a contrastive tillage site in a similar topography condition in Loess Plateau, China. The purpose was to explore the dynamics of soil organic carbon (SOC) and dissolved organic carbon (DOC) in *R. pseudoacacia* forests. On average depths of 20, 40, and 60 cm, SOC, active organic carbon (AOC), and DOC gradually increase with increased forest age. After forest restoration, the AOC/SOC ratio and resistant organic carbon/SOC ratio increase, whereas the slow organic carbon/SOC ratio decreases. The soil solutions in the subsoil layer have low DOC:DON ratio and high UV absorption at 280 nm. At 40 and 60 cm, the depth distribution is indicated as special low values for DOC concentration in the C99 site (10a site), as well as for soil water content, SOC, and AOC in the 25a forest site. Our results provide evidence that during forest restoration, SOC does not consistently increase linearly. The change points of different SOC proportions and DOC concentrations at various depths are not same, i.e., asynchronous changes exist.

Keywords: depth distribution, soil organic carbon dynamics, active organic carbon (AOC), UV absorption at 280 nm.

1. Introduction

Soils have the potential for carbon (C) release or sequestration. Hence, soil organic carbon (SOC) is related to atmospheric CO₂ levels, which can be affected by vegetation cover and land management (Lal, 2004). When land is converted from one cover type to another, research on SOC dynamics is valuable in improving our comprehension and increasing our predictive capability over both short- and long-term scales (Post and Kwon, 2000). Land cover changes affect litters, plant root, soil fauna, soil microorganisms, and soil condition, which can drastically alter the soil carbon stock. Generally, the transformation of cultivated soil to forest ecosystem can realize additional soil carbon sequestering. The concomitant development of soil carbon and vegetation in primary and secondary chronosequences was documented by Bush *et al.* (2008), among others. SOC increases by up to 52% after a change from crop to secondary forest (Guo *et al.*, 2002). Forests with nitrogen-fixing trees may also allocate a higher fraction of their net primary production to soil (e.g., Bush, 2008; Kiser *et al.*, 2009). However, the results of soil carbon stock change in forest chronosequence are likely to be different. The increasing trend of soil carbon is not always represented in the entire forest chronosequence or at each depth. Zhou *et al.* (2006) found that pioneer and transition forests had limited effects on SOC accumulation even if the aboveground biomass rapidly accumulates and reaches a high level. In the hilly Loess Plateau of China, Wang *et al.* (2010) indicated that SOC contents are significant for different land covers in a semi-arid climate only at 5 cm soil surface depth. Cropland transformation to immature woodland has no significant effect on SOC.

SOC is greatly susceptible to environment change, especially to vegetation cover changes (Wang *et al.*, 2011). Large amounts of worldwide soil carbon data

need to be analyzed to explore highly inherent natural variability in worldwide soils and carbon loss rules under different land cover types and change trajectories (Lal, 2009). The mechanism of SOC dynamics in forest chronosequences needs to be determined.

In forest, grass, and shrub vegetation restoration, DOC leaching plays an important role in SOC at different depths, which is a crucial function in the belowground carbon cycle. The largest increase in the flux of dissolved organic matter (DOM) occurs when percolating water passes through the forest floor (Michalzik *et al.*, 2001). DOM leaching from organic layers to mineral soils and DOM sorption onto mineral surfaces are very likely to be important mechanisms in stabilizing soil organic matter (Kalbitz and Kaiser, 2008). Therefore, both SOC and DOC should be calculated and measured in predicting ecosystem response to global change, especially for depth distribution.

The Loess Plateau in China covers approximately 58×10^4 km². In the 1970s, the Chinese government began an extensive tree plantation. Since then, the Zhifanggou drainage area of the Ansai country of Shaanxi province has undergone remarkable land use change. The conversion of degraded lands into forests and grasslands was essentially promoted by the state-funded project, Grain-for-Green, in 1999. The eco-environment was improved under government regulations of reforestation/planting grass (Fu *et al.*, 2010). The restored forest species included *Platycladus orientalis*, *Quercus variabilis*, *Robinia pseudoacacia*, and *Pinus tabulaeformis*, among others. In the Zhifanggou drainage basin, one of the main restored species, *R. pseudoacacia*, was planted extensively as a kind of pioneer forest plant because of its high adaptability. Restored *R. pseudoacacia* forests of different ages were mixed with one another at patch scales in the Zhifanggou drainage basin, enabling the selection

of suitable sites in a small field. The present study used a special comparison method to (1) estimate SOC dynamics in a period of 35 years for *R. pseudoacacia* forest restoration in a similar topographic condition, (2) investigate the mechanism of SOC and DOC dynamics during the development of forest vegetation, and (3) explore the depth distribution of SOC and its potential change mechanism in forest restoration.

2. Materials and Methods

2.1 Field site and sampling regime

The study area is the Zhifanggou drainage basin ($109^{\circ}13'46''$ – $109^{\circ}16'03''$ E, $36^{\circ}46'42''$ – $36^{\circ}46'28''$

N), located in the middle part of the Loess plateau in northern Shaanxi Province, P. R. China (Figure 1). Due to significant topographic variations within the Loess hills and gully landforms, only a small area of $275\text{ m} \times 212.5\text{ m}$ is covered to ensure that all selected sites had similar topographic conditions. Four *R. pseudoacacia* forest sites and one tillage site were selected (Table 1). The region mainly has a semi-arid climate of warm temperate zone, with an annual accumulative radiation of 493 kJ cm^{-2} . The average annual temperature is $8.8\text{ }^{\circ}\text{C}$ and the mean annual precipitation is 549.1 mm . The soil in the study area is fine silt to silt in texture and weakly resistant to erosion.




Figure 1. Location of the study sites with a 2 m contour lines.

Table 1. Summary of land-use history, geographic condition, vegetation and planting details of five study sites.

Sample sites	Stand age (years)	Acreage (m ²)	Old land use	Aspect	Slope (°)	Crown density	Total litters (g m ⁻²)	Trunk diameter (cm)	Tree height (m)
Tillage	∞	1220	—	W	20.2	—	—	—	—
C99	10	1581	Tillage	SWW	27.3	0.9-0.95	348.3	11.4	11-12
C84	25	913	Tillage	SWW	21.4	0.5-0.6	262.4	12.8	15-20
C78	31	1344	Tillage	SWW	21.4	0.75-0.8	362.9	23.8	15-20
C74	35	486	Tillage	SWW	18.3	0.85-0.9	298.6	30.1	>20

In 1973, the Zhifanggou basin was selected as an experimental site by the Key Technologies Research and Development Program on the comprehensive management of the Loess Plateau. Partial farmlands were reverted into forests and grasslands in batches. Four kinds of *R. pseudoacacia* forests were respectively planted in 1999 (10a or C99), 1984 (25a or C84), 1978 (31a or C78), and 1974 (35a or C74). The forests herbs include *Arundinella hirta*, *Roegneria kamoji*, *Viola yedoensis*, *Aneurolepidium dasystachys*, *Artemisia japonica*, *Carex rigescens*, *Agropyron cristatum*, etc. The tillage study site has existed since 1974 and is less affected by chemical fertilizer. When cropland in this area was gradually restored to forest, grassland, or shrub, the tillage site was kept but the corns, soybean, and millet were rotated. Four forest sites and the tillage site locate in 200-500 m east to Shiyaoxian village, where all restored forest fields were distributed to villagers in order to manage effectively. According to old villager's introduction, in history the study area had been utilized as farmland by serious food demand until 1973.

An average of nine sections was selected in each study site. Samples from depths of 20 ± 2 , 40 ± 2 , and 60 ± 2 cm were obtained using a soil drill and evenly mixed with three horizontal samples, respectively. We

enclosed these samples in a plastic bag and stored them in the laboratory at 2 °C. Three 100 cm-deep holes were randomly selected and dug in each study site. Samples were collected from depths of 5–10, 30–35, 60–65, and 90–95 cm, and stored in 50 mL aluminum boxes to determine the soil water content and soil bulk density.

2.2 Sampling and analysis

The moisture samples in aluminum boxes were weighed wet, dried at 105 °C for 48 h, and weighed again to determine the bulk density and gravimetric soil water content. After air drying soil samples for 3 days, litters were carefully picked out, crushed to pass through a 1 mm sieve, and stored at 2 °C until analysis.

In 1989, Parton (1989) divided SOC into active (AOC), slow (LOC), and resistant C pool (ROC) based on the mean residence times. In the present study, SOC was determined using the routine potassium dichromate oxidation-outer heating method. The 333 mM KMnO₄ chemical oxidation method was used for AOC determination (Conteh *et al.*, 1997). ROC was determined by the acid hydrolysis method (Collins *et al.*, 2000). About 1 g of soil samples were hydrolyzed with 20 mL of 6 M HCl in sealed Pyrex tubes at 105 °C for 18 h. The hydrolysate was then discarded.

After washing unhydrolyzed residues with deionized water by repeated centrifugations and decantations, the samples were transferred to pre-weighed vials and dried at 60 °C to constant weight. Subsequently, SOC was analyzed. LOC was calculated by difference between SOC and ROC ($LOC = SOC - ROC$).

DOM was collected using a soil-to-solution (distilled water) ratio of 1:5, shaken for 1 h, and centrifuged (2000 rpm for 10 min). The solution was collected, filtered through 0.45 µm cellulose acetate filters, and stored in a frozen state until analysis. DOC was determined with a TOC Analyzer (Phoenix 8000). Total nitrogen (TN) in the solution was determined using a photometer after persulfate digestion. Inorganic nitrogen was determined using a continuous flow analyzer (AA3). Dissolved organic nitrogen (DON) was determined by the difference between TN and dissolved inorganic nitrogen (DIN), i.e., $DON = TN - DIN$. The UV absorbance at 280 nm was measured (adjustive solution of 10 mg L⁻¹ DOC, calculated by dividing measured UV absorption at 280 nm) to estimate aromaticity (Zsolnay *et al.*, 1999). All samples were determined with three replicates.

2.3 Data handling and statistics

All figures and most data were made using Excel 2003 and SPSS 16.0 software. One-way ANOVA was used to examine the effects of the vegetation restoration type on the concentrations and stocks of soil carbon. If significant effects were observed by ANOVA, a least significant difference [LSD (0.05)] test was used.

3. Results

3.1 Soil water content and bulk density

A comparison of the tillage and forest sites revealed that their soil bulk density had big difference ($p = 0.05$) at 5–10 cm, but did not exhibit coherent diversity in subsoil (Table 2). Soil water content was obviously higher in the tillage site than in the forest sites. With increased forest restoration age, the soil water content approximately increased. Soil water content in the C84 site showed no obvious change from 60–95 cm. Water content was lower in C78 than in C84. With increased depth, the soil water contents in all study sites increased, except for C84.

Table 2. Bulk density and water content of 100 cm soils from the five study sites.

variables	sites	5-10cm	30-35cm	60-65cm	90-95cm
Bulk density (g cm ⁻³)	Tillage	1.24±0.16 ^b	1.23±0.11 ^b	1.22±0.20 ^b	1.36±0.10 ^a
	C99	1.05±0.34 ^c	1.16±0.20 ^c	1.23±0.30 ^b 0154	1.17±0.15 ^c .111
	C84	1.43±0.29 ^a	1.44±0.15 ^a	1.29±0.21 ^{ab}	1.16±0.20 ^c
	C78	1.38±0.54 ^a	1.16±0.25 ^c	1.34±0.17 ^a	1.38±0.15 ^a
	C74	1.38±0.33 ^a	1.26±0.15 ^b	1.17±0.20 ^c	1.27±0.11 ^b
Water content (%)	Tillage	4.74±0.11 ^a	6.98±0.36 ^a	7.60±2.30 ^a	8.33±2.01 ^a 2
	C99	2.32±0.27 ^b	3.36±0.21 ^c	4.33±1.29 ^c	4.86±0.56 ^c
	C84	2.81±1.22 ^b	3.12±0.36 ^{cd}	3.18±0.64 ^d	2.98±1.00 ^d
	C78	2.55±0.75 ^b	2.90±1.01 ^d	3.36±0.99 ^d	5.09±1.53 ^{bc}
	C74	4.07±0.65 ^a	4.47±0.13 ^b	5.33±1.64 ^b	5.68±2.06 ^b

Values are means±standard errors of n=3. Significant differences between sites are indicated by superscript letters (LSD; $p \leq 0.05$).

3.2 Changes in SOC, AOC, and DOC

For the three depths, the change scope of SOC was smaller in the tillage site than in the forest sites

(Table 3). The average SOC of the three depths increased with increased forest age.

Table 3. SOC, AOC and DOC quantity of five study sites.

Variables	Depths	Tillage	C99	C84	C78	C74
SOC (g kg ⁻¹)	20cm	2.23±0.13 ^b	2.58±0.14 ^{ab}	3.32±0.40 ^{ab}	3.75±0.30 ^{ab}	6.65±1.66 ^a
	40cm	1.73±0.05 ^b	1.68±0.07 ^{bc}	1.37±0.07 ^c	1.97±0.03 ^b	3.31±0.12 ^a
	60cm	1.86±0.04 ^b	1.38±0.05 ^c	1.14±0.02 ^c	1.99±0.12 ^b	3.05±0.07 ^a
	average	1.92±0.03 ^b	1.81±0.10 ^b	1.82±0.65 ^b	2.46±0.56 ^{ab}	4.13±1.03 ^a
AOC (g kg ⁻¹)	20cm	0.09±0.01 ^a	0.19±0.03 ^a	0.23±0.05 ^a	0.25±0.02 ^a	0.49±0.16 ^a
	40cm	0.04±0.00 ^d	0.07±0.01 ^c	0.04±0.00 ^{cd}	0.11±0.01 ^b	0.17±0.01 ^a
	60cm	0.03±0.00 ^c	0.03±0.00 ^c	0.02±0.01 ^c	0.06±0.00 ^b	0.11±0.00 ^a
	average	0.05±0.00 ^b	0.09±0.01 ^b	0.09±0.011 ^b	0.13±0.01 ^{ab}	0.24±0.02 ^a
DOC (mg kg ⁻¹)	20cm	14.57±2.32 ^{bc}	16.50±5.36 ^b	12.80±6.11 ^c	20.86±1.40 ^a	25.47±7.56 ^a
	40cm	13.26±3.26 ^c	2.76±0.54 ^d	12.64±8.92 ^c	18.50±0.20 ^b	24.40±2.17 ^a
	60cm	16.01±1.03 ^a	2.56±0.85 ^b	13.82±0.32 ^a	16.15±5.45 ^a	14.32±3.16 ^a
	average	14.61±5.61 ^b	7.27±1.23 ^c	13.09±5.13 ^{bc}	18.50±6.00 ^{ab}	21.40±4.26 ^a

Values are means±standard errors, n=9 besides average. Averages are calculated with 27 data (n=27) of three depths in each site. Significant differences between sites are indicated by superscript letters (LSD test; $p \leq 0.05$).

Compared with tillage, the average SOC values in the 10a and 25a forest sites decreased, increased to the original level (value of the tillage site) in the 31st year of restoration, and increased continually thereafter. SOC increased with increased age at 20 cm in the C99, C84, C78, and C74 forest sites. However, SOC decreased at 40 and 60 cm in C99 and C84. On average, the SOC at three depths and at 20 cm had more remarkable diversity degrees between the forest and tillage sites. SOC in C74 (35a) was significantly

higher than that in the tillage site (distinct diversity degree, $p = 0.05$) in each depth.

With increased age, AOC increased at 20 cm in the four forest sites. However, the AOC values at 40 and 60 cm in C84 (25 a) were lower than that in the tillage site. LOC in the forest sites obviously increased compared with that in the tillage site. Given the low AOC value at 40 and 60 cm, AOC in C84 was lower than that in C99, but still 84.0% higher than that in the tillage site.

The DOC values at 40 and 60 cm in C99 were lower, but those at 20 cm were higher than those in the tillage site. DOC at 20 cm gradually increased with increased age, except for DOC in C84, which was 3.7.1 mg kg⁻¹ lower than that in C99. DOC values at 40 and 60 cm in C99 were lower than that in the tillage site. From 10 years to 25 years, the DOC values at 40 and 60 cm increased gradually and approached that of the tillage site after 25 years.

3.3 SOC proportion

In all four forest sites, AOC/SOC and ROC/SOC were higher but LOC/SOC was lower than those of the tillage site (Table 4). The AOC/SOC ratio in C84 (25a) was lower than in C99 (10a) at the three depths, but showed no obvious difference between C74 (35a) and C78 (31a). With increased depth, the AOC/SOC ratio decreased gradually. In all four forest sites, AOC/SOC at 60 cm was 44.9%–69.4% lower than that at 20 cm, compared with the 64.1% for the tillage site.

Table 4. Contrast of three SOC proportions in five study sites.

Depths	Variables	Tillage	C99(10a)	C84(25a)	C78(31a)	C74(35a)
20cm	AOC/SOC	0.04±0.00 ^b	0.07±0.01 ^a	0.06±0.01 ^a	0.07±0.00 ^a	0.07±0.01 ^a
	LOC/SOC	0.68±0.01 ^a	0.50±0.05 ^a	0.49±0.05 ^a	0.64±0.01 ^a	0.50±0.04 ^a
	ROC/SOC	0.29±0.02 ^a	0.43±0.05 ^a	0.45±0.04 ^a	0.29±0.01 ^a	0.43±0.03 ^a
40cm	AOC/SOC	0.02±0.00 ^b	0.04±0.01 ^{ab}	0.03±0.00 ^b	0.05±0.00 ^a	0.05±0.00 ^a
	LOC/SOC	0.70±0.01 ^a	0.65±0.03 ^{ab}	0.59±0.03 ^{ab}	0.57±0.03 ^b	0.606±0.03 ^{ab}
	ROC/SOC	0.28±0.01 ^a	0.31±0.03 ^a	0.38±0.03 ^a	0.38±0.03 ^a	0.35±0.03 ^a
60cm	AOC/SOC	0.01±0.00 ^c	0.02±0.00 ^b	0.02±0.00 ^{bc}	0.03±0.00 ^{ab}	0.04±0.00 ^a
	LOC/SOC	0.74±0.03 ^a	0.65±0.03 ^{ab}	0.70±0.03 ^{ab}	0.53±0.01 ^b	0.62±0.02 ^b
	ROC/SOC	0.24±0.03 ^b	0.33±0.03 ^{ab}	0.28±0.03 ^b	0.44±0.01 ^a	0.35±0.02 ^{ab}

Values are means±standard errors, n=9 besides average. Significant differences between sites are indicated by superscript letters (Tukey-Kramer test; $p \leq 0.05$).

3.4 DOC/DON ratio and UV absorption at 280 nm of DOM

With increased depth, the DOC/DON ratio in all study sites gradually increased (Figure 2). The DOC/DON ratio at 20 cm of C78 was close to that

of the tillage site, whereas those of the other forest sites decreased to different extents. The lowest value appeared in C84, accounting for 29.9% of the tillage site. At 40 and 60 cm, the DOC/DON values decreased. At 60 cm, DOC/DON decreased to 36.1%–58.3%.




Figure 2. DOC quality changes in soil solution of three depths. Means and standard errors of nine replicates.

UV absorbance decreased with increased depth in each site. At 20 cm, the UV absorbance of C78 accounts for 39.1% of that of C84. The UV absorbance index at 20 and 40 cm was higher in each forest site than that in the tillage site, and increased with increased restoration age. The UV absorbance index at 60 cm in C99 and C78 approached that of the tillage site, but was higher in C84 and C74 than in the tillage site.

4. Discussion

4.1 Soil water content

Soil water content is affected by both vegetation cover on the soil surface and plant roots in subsoil. Soils of forest vegetation in the Loess plateau have lower water content than those in tillage sites because of root

absorption (Zhao, 2005). Vegetation restoration changes the litter density, root quantity, and ultimately, soil conditions such as pH, temperature, moisture, microorganisms, etc., which are indicated as significant factors affecting soil carbon turnover (Lal, 2005). The results of the evaporation and herbage root absorption at 20 cm, as well as the more powerful arbor root absorption at 40 and 60 cm show that the water content at three depths did not exhibit the same change trend with increased age. With the development of forest vegetation, the soil water content decreases in subsoil and soil surface, but increases overall. The soil water content changes the conditions of the organic carbon pool and affects its turnover rate, deposited form, and SOC proportion. In semi-arid areas, high soil water contents increase SOC concentration with high humification rates. Hence, the depth distribution of water content can at least partly affect the SOC proportion and concentration.

4.2 SOC concentration and proportion changes during 35 years forest restoration

Litters are the most important sources of SOC in the forest ecosystem (Huang *et al.*, 2011). However, the carbon input on surface soil is higher than in deeper soil layers. When tillage was converted into forest, litters accumulated gradually on the soil surface and the depth distribution of SOC was enhanced. Compared with tillage, the depth distribution of SOC in the 10a to 35a forests acutely increased at 20 cm due to litter accumulation. Subsoil organic carbon is mainly generated from root exudates, leaches DOC, fragmentation, and particle SOC movement, which produce limited C input. However, subsoil layers in the tillage site can intercept more organic carbon than those in the forest by ploughing and fertilization, which contribute to the decrease in subsoil organic carbon in young forest restoration.

In general, the amount of SOC is greater in forests than in arable soils (Vesterdal *et al.*, 2002). For example, in tropical areas, SOC increases by 29% when agricultural land is afforested (Axel *et al.*, 2011). However, a land-use change from arable to forest does not always increase the sequestration of soil organic matter (Vesterdal *et al.*, 2002). In our study, although the average SOC in the four forest sites increased gradually, those in the C99 and C84 forest sites were less than that in the tillage site. SOC at 40 and 60 cm showed different changes compared with that at 20 cm, suggesting the diversity of SOC turnover mechanisms between the soil surface and subsoil. Fontaine *et al.* (2007) found that adding fresh SOC can accelerate old SOC decomposition. After forest restoration, SOC in subsoil was changed by some factors, such as litter fragmentation, SOC composition, and root density, different from that on the surface soil, indicating an asynchronous state. In the 25a forest site, soil water content, SOC, and AOC congruously displayed lower values at 40 and 60 cm. During forest restoration, SOC in subsoil initially increased due to litter fragmentation, which led to a slight increase in the 10a site, and then decreased due to old SOC decomposition and root absorption. Our study proved that the fluctuation of SOC was represented only in the first 25 years of forest restoration. The mechanism of the “decrease factor” such as old SOC decomposition in response to the “increase factor” such as litter fragmentation still needs to be explored.

By the nitrogen fixation function of *R. pseudoacacia*, litters on the soil surface with lower C/N ratios led to fast decomposition times, accounting for high AOC quantity (MA *et al.*, 2007). In all four forest sites, the AOC/SOC and ROC/SOC ratios were higher than those in the tillage site. This finding suggested that after forest restoration from tillage, AOC ratio and ROC ratio increased, and LOC ratio was consumed. Hence, forest vegetation may help restore carbon in soil. An

asynchronous state was also suggested by the SOC proportions. In our study, AOC was regarded had the highest biochemical quality within the shortest residence time, and played an important role in plant nutrition. ROC is an inert compound with stable physical and chemical qualities. LOC can be deemed as a recalcitrant compound with the longest residence time (Paul *et al.*, 1999). Kazumi (2011) found that young, fast-MRT soil carbon is decomposed in the upper A horizon, and old, stable soil carbon is decomposed in the lower A horizon in the forest. In the present study, the AOC/SOC ratio in all four forest sites decreased gradually with increased depth. The SOC dynamics and compositions differed between the soil surface and subsoil, leading to different changes during forest restoration.

4.3 DOC changes in forest sites

DOC in soil solution is generated from litter leachates, root exudates, and microbial degradation products (Zsolnay, 1996). In the forest ecosystem, litters on the forest floor gradually accumulates, microbial degradation products increase due to high AOC values with short residence times, and root density in subsoil increase. Consequently, the average DOC increases with increased age.

The DOC values at 40 and 60 cm in 10a forest were lower than in the other study sites, contributing to lower average DOC values in 10a forest than in the tillage site. Theoretically, both DOC leaching from soil surface and root exudates can lead to the low depth distribution of the study sites, except the 10a forest site. In other words, they decrease the DOC difference between the soil surface and subsoil layer. As important sources, the degradation products of litters and AOC on the soil surface hardly leach through the subsoil layer (Matthias, 2009), especially when AOC on the soil surface is low in the beginning of forest restoration. For young forests, the plant root density and microbe quan-

tity have low values, low root exudates, and microbial degradation, leading to low DOC values in the subsoil layer in the 10 a site, although the AOC value was higher than that in the 25 a forest site. With increased root density and microbe quantity, subsoil SOC and AOC in 25a forest had the lowest values, whereas DOC increased compared with 10a forest.

UV absorbance and DOC: DON were utilized to evaluate DOM property and composition (Traina *et al.*, 1990). The specific UV absorbance at 280 nm was considered as a measure of aromaticity (Chin *et al.*, 1994), and DOC:DON was considered as a proxy for the quality of the DOM. Increased DOC:DON indicates reduced bioavailability (e.g., Brian *et al.*, 2009). In our study, DOC:DON at 20 cm in the forest sites showed no obvious difference with those in the tillage site; however, those at 60 cm were higher than those in the tillage site. Stephan (2011) found the throughfall contribution to the overall element fluxes was higher for DON than for DOC. Lower DOC/DON values in forest than in tillage sites reflects that after forest restoration, soil solution movement is more active, and/or more recalcitrant DOM is removed from the solution (Brian *et al.*, 2009). On the other hand, forest restoration leads to changes in SOC proportions, such as increased AOC and ROC ratio, as well as decreased LOC, suggesting biodegradation action. A part of DOC is released from SOC biodegradation. DOC in forest sites mainly consist of litter-derived DOC and SOC-derived DOC, providing lower DOC/DON values and higher aromaticity compositions than in tillage sites.

5. Conclusions

When tillage was reverted into forest, SOC exhibited a total, nonlinear, absolute increase, suggesting asynchronous changes. During *R. pseudoacaci* forest restoration, the difference between the soil surface and

subsoil led to the depth distribution of SOC and AOC. The 25a forest represented a special phase wherein soil water content, SOC, and AOC exhibited low values in the subsoil layer. Apparently, there was a natural C-input course during forest restoration.

After forest restoration, the total SOC and AOC increased and ROC increased, suggesting an internal conversion process among the SOC proportions. In forest ecosystems, the LOC concentration decreases and the degradation and transformation among AOC, LOC, and ROC changed. However, whether the changes benefit the soil carbon input in the long run is still unclear.

In the beginning of forest restoration, DOC concentration at 40 and 60 cm sharply decreased, suggesting obvious depth distribution. However, the depth distribution of DOC for mature forests was not obvious. DOC does not perfectly indicate the same changes as those of SOC proportions, and more factors control the DOC concentration.

Acknowledgments

We thank Jinyi Wang from Northwest A&F University for academic support, who improved the presentation of the manuscript. This work was supported by Knowledge Innovation Project of The Chinese Academy of Sciences □KZCX2-XB2-05-01 □

References

- Axel, D., Jens, S., Annette, F. 2011. Impact of tropical land-use change on soil organic carbon stocks- a meta-analysis. *Global Change Biology*. 17, 1658-670.
- Brian, D.S., Robert, B. H., Thomas, A.T., *et al.* 2009. Changes in dissolved organic matter with depth suggest the potential for postharvest organic matter retention to increase subsurface soil carbon pools. *Forest Ecology and Management*. 258, 2347-352
- Bush, J.K. 2008. Soil nitrogen and carbon after twenty years of riparian forest development. *Soil Science Society of America Journal*. 72, 815-822.
- Chin, Y. P., Alken, G., O'Loughlin, E. 1994. Molecular weight, polydispersity, and spectroscopic properties of aquatic humic substances. *Environmental Science and Technology*. 28, 1853-1858.
- Collins, H.P., Elliott, E.T., Pustian, K. *et al.* 2000. Soil carbon pools and fluxes in long-term corn belt agroecosystems. *Soil Biology and Biochemistry*. 32, 157-168.
- Conteh, A., Lefroy, R.D.B., Blair, G.J. 1997. Dynamics of organic matter in soils as determined by variations in ¹³C/¹²C isotopic ratios and fractionation by ease of oxidation. *Australian Journal of Soil Research*. 35, 881-890.
- Fontaine, S., Barot, S., Barre, P., Bdioui, N., Mary, B., Rumpel, C. 2007. Stability of organic carbon in deep soil layers controlled by fresh carbon supply. *Nature*. 450, 277-280.
- Fu, X.L., Shao, M.A., Wei, X.R. *et al.* 2010. Soil organic carbon and total nitrogen as affected by vegetation types in Northern Loess Plateau of China. *Geoderma*. 155, 31-35.
- Guo, L.B., Gifford, R.M., 2002. Soil carbon stocks and land use change: a meta analysis. *Glob. Change Biol.* 8, 345-360.
- Huang, Y.H., Li, Y.L., Xiao, Y. *et al.* 2011. Controls of litter quality on the carbon sink in soils through partitioning the products of decomposing litter in a forest succession series in South China. *Forest ecology and management*. 261, 1170-1177.
- Kalbitz, K., Kaiser, K. 2008. Contribution of dissolved organic matter to carbon storage in forest mineral soils. *Journal of Plant Nutrition and Soil Science*. 171, 52-60.

- Kazumi, K., Yosuke, M., Kenji, T. *et al.* 2011. A study of the carbon dynamics of Japanese grassland and forest using ^{14}C and ^{13}C Nuclear Instruments and Methods in Physics Research. B. 268, 1106-1109
- Kiser, L.C., Kelly, J.M., Mays, P.A. 2009. Changes in forest soil carbon and nitrogen after a thirty-year interval. *Soil Science Society of America Journal*. 73, 647-653.
- Lal, R. 2004. Soil carbon sequestration impacts on global climate change and food security. *Science*. 304, 1623-1627.
- Lal, R. 2005. Forest soils and carbon sequestration. *Forest ecology and management* 220, 242-258.
- Ma, Y.H., Guo, S.L., Yang, Y.L. *et al.* 2007. Influence of Vegetation Types on Soil Organic C at Yangou Catchment in the Loess Hilly-gully Region *Journal of natural resources (in Chinese)* 22(1), 97-106
- Michalzik, B., Kalbitz, K., Park, J.H., Solinger, S., Matzner, E. 2001. Fluxes and concentrations of dissolved organic carbon and nitrogen-a synthesis for temperate forests. *Biogeochemistry*. 52, 173-205.
- Post, W.M., Kwon, K.C. 2000. Soil carbon sequestration and land-use change: processes and potential. *Glob. Change Biol.* 6, 317-327.
- Parton, W.J., Sandford Jr, R.L., Sanchez, P.A., Stewart, J.W.B. 1989. Modeling soil organicmatter dynamics in tropical soils. *Soil Science Society of America Journal*. 39, 153-171.
- Paul, E.A.D., Harris, H.P., Collins, U.S., Robertson, G.P. 1999. Evolution of CO_2 and soil carbon dynamics in biologically managed, rowcrop agroecosystems. *Applied Soil Ecology*. 11, 53-65.
- Traina, S.J., Novak, J., Smeck, N.E. 1990. An ultraviolet absorbance method of estimation the percent aromatic carbon content of humic acids. *Journal of Environmental Quality*. 19, 151-153.
- Vesterdal, L., Ritter, E., Gundersen, P. 2002. Change in soil organic carbon following afforestation of former arable land. *Forest Ecology Management*. 169, 137-147.
- Wang, Y.F., Fu, B.J., Lü, Y.H., Song, C.J., Luan, Y. 2010. Local-scale spatial variability of soil organic carbon and its stock in the hilly area of the Loess Plateau, China. *Quatern Res.* 73, 70-76.
- Wang, Y.F., Fu, B.J., lv, Y.H. *et al.* 2011. Effects of vegetation restoration on soil organic carbon sequestration at multiple scales in semi-arid Loess Plateau, China. *Catena*. 85, 58-66.
- Zhao, J.B., Du, J., Zhou, Q. *et al.* 2005. Dry layer of soil below artificial forest near XianYang in Shaanxi. *Scientia geographica sinica*. 25(3), 322-329. (in Chinese)
- Zhou, G.Y., Zhou, C.Y., Liu, S.G. *et al.* 2006. Below-ground carbon balance and carbon accumulation rate in the successional series of monsoon evergreen broad-leaved forest. *Science in China Series D: Earth Sciences*. 49, 311-321.
- Zsolnay, A. 1996. Dissolved humus in soil waters. In: Piccola, A. (Ed.). *Humic Substances in Terrestrial Ecosystems*. Elsevier. Amsterdam pp, 171-223.
- Zsolnay, A., Baigar, E., Jimenez, M. *et al.* 1999. Differentiating with fluorescence spectroscopy the sources of dissolved organic matter in soils subjected to drying. *Chemosphere*. 8, 45-50