

Short communication

Utility of five SSR markers for genetic diversity and paternity exclusion analysis in the Patagonian toothfish

Cristian Araneda¹, Natalia Lam^{1,2}, Patricia Iturra², Felipe Jilberto¹
Valentina Cordova² & Pablo Gallardo³

¹Departamento de Producción Animal, Facultad de Ciencias Agronómicas
Universidad de Chile, Santiago, Chile

²Programa Genética Humana, ICBM, Facultad de Medicina
Universidad de Chile, Santiago, Chile

³Escuela de Ciencias y Tecnologías Agrícolas y Acuícolas, Facultad de Ciencias
Universidad de Magallanes, Punta Arenas, Chile

Corresponding author: Pablo Gallardo (pablo.gallardo@umag.cl)

ABSTRACT. The Patagonian toothfish or Chilean sea bass (*Dissostichus eleginoides*), found in the Southern Ocean surrounding Antarctica, is an important fishery species for Chile. This high-value species is regarded as overfished, making it an attractive target for aquaculture. When developing a reproduction program for any aquaculture species, it is important to implement genetic tools to evaluate diversity, inbreeding, and parentage. We calculated genetic diversity and paternity/maternity exclusion probabilities based on five commonly-used microsatellite *loci* in a natural population of Patagonian toothfish from southern Chile (n = 34) in order to evaluate the potential utility of these five markers in stock management. The observed number of alleles per *locus* (N_a) and observed heterozygosities (H_o) are within range as described by studies performed in other sub-Antarctic regions. All five *loci* were strongly polymorphic, with $H_o > 0.6$ and $N_a > 7$, and paternity exclusion probabilities were high ($PE > 0.99$), indicating the potential utility of these *loci* in paternity/maternity exclusion analyses of Patagonian toothfish.

Keywords: *Dissostichus eleginoides*, Chilean sea bass, heterozygosity, microsatellites, genetic diversity.

The Patagonian toothfish (*Dissostichus eleginoides*), also marketed as “Chilean sea bass,” is an important commercial fishery species found in the Southern Ocean surrounding Antarctica. It has been regarded as overfished since the 1990s (Marko *et al.*, 2011; Reyes *et al.*, 2012). The species is a benthopelagic fish with a long lifespan and is widely distributed along the southern Chilean coast, Patagonian shelf, and sub-Antarctic islands and seamounts at depths of 1000-2000 m (Appleyard *et al.*, 2002). Commercial fishing of the Patagonian toothfish is on the rise due to its market value (which is higher than that of salmon, for instance), and international regulations govern sustainable fishery of the species (Marko *et al.*, 2011). Therefore, the Patagonian toothfish is an attractive target for Chilean aquaculture (Reyes *et al.*, 2012). Chilean government agencies currently provide strong incentives and funding for aquaculture programs with

new marine species. In Chilean Patagonia, aquaculture research projects are establishing *Dissostichus eleginoides* broodstock as a basis for developing culture technology. This research is focused on obtaining high-quality juveniles, addressing issues such as nutrition and genetics, and maximizing maturation and reproductive performance. In this context, genetic tools are needed to support reproduction programs in assessing genetic diversity, inbreeding, and parentage. Short-sequence repeats (SSR), or microsatellites, with high levels of polymorphism have been the most useful molecular markers for evaluating marine fish populations. We tested five microsatellite markers (cmrDe2, cmrDe4, cmrDe9, cmrDe13, and cmrDe30) (Reilly & Ward, 1999) in an offshore Chilean population. These markers had been commonly used in genetic diversity studies to evaluate the distribution and population structure of the Patagonian toothfish in the

Southern Ocean, Antarctic Islands, Patagonian shelf (Falkland/Malvinas Islands), and Atlantic, West and Western Indian Ocean sectors (Smith & McVeagh, 2000; Appleyard *et al.*, 2002, 2004; Shaw *et al.*, 2004; Rogers *et al.*, 2006). Our main objectives were to evaluate genetic diversity and estimate paternity/maternity exclusion probabilities, in order to evaluate the potential utility of these five markers in broodstock management.

We studied 34 fishes, collected in September-October of 2011 from two sampling points in Chilean Patagonia in front of Trinidad Gulf (Magallanes and Chilean Antarctic region): $n = 19$ from point one ($49^{\circ}29'41''\text{S}$, $76^{\circ}26'17''\text{S}$) and $n = 15$ from point two ($50^{\circ}15'05''\text{S}$, $76^{\circ}20'15''\text{W}$). Genomic DNA was extracted using the phenol-chloroform method from fin-clip samples (Taggart *et al.*, 1992). Microsatellite analysis was performed using five markers developed for this species (*cmrDe2*, *cmrDe4*, *cmrDe9*, *cmrDe13*, and *cmrDe30*) (Reilly & Ward, 1999) and the M13-tailed primer method (Schuelke, 2000) (Table 1). PCR amplifications were carried out as 15- μL reactions containing 10x PCR buffer (10 mM Tris-HCl, 50 mM KCl, 1 mM EDTA, Triton X-100 0.1%), 2.5 mM MgCl_2 , 0.2 mM dNTPs, 8 pmol reverse primer, 8 pmol fluorescent-labeled universal M13 primer, 2 pmol forward primer, 1 U Taq polymerase (Biotools[®]), and 2- μL DNA samples. The thermal profile was 94°C for 2 min, followed by 30 cycles at 94°C (30 s), specific marker T_m (Table 1) (45 s), 72°C (45 s), 8 additional cycles at 94°C (10 s), 53°C (45 s) to bind M13-labeled primer, and 72°C (45 s) with a final 10-min extension step at 72°C .

Different amplicons labeled with different fluorescent dyes were pooled in one reaction tube and run with a GeneScan[™] 500 LIZ[™] dye-size standard (Applied Biosystems[®]) in an automatic sequencer at external laboratories. Genotypes were identified using GeneMarker[®] software (SoftGenetics[®]). Genetic diversity was assessed according to observed (H_o) and expected (H_e) heterozygosities, number of observed alleles by locus (N_a), polymorphism information content (PIC), and paternity/maternity exclusion probabilities (PE), with Cervus 3.0 software (Araneda *et al.*, 2004; Kalinowski *et al.*, 2007). The presence of null alleles was assessed with six different estimators (Dempster *et al.*, 1977; Chakraborty *et al.*, 1992; Brookfield, 1996; Van Oosterhout *et al.*, 2004; Kalinowski *et al.*, 2006), using Micro-Checker 2.2.3 (Van Oosterhout *et al.*, 2004), Cervus 3.0 (Kalinowski *et al.*, 2007) and FreeNA (Chapuis & Estoup, 2007). Genepop 4.2 (Rousset, 2008) was used to evaluate deviation from Hardy-Weinberg expectations along with pairwise F_{ST} to evaluate population differentiation between the two collection sites.

Genetic differentiation did not vary significantly between the two sampling points ($F_{ST} = -0.002$; $P > 0.05$), possibly due to their geographical proximity. Therefore, in subsequent analyses, all 34 samples were used together and considered to belong to the same biologic population.

N_a ranged from 8 (*cmrDe13*) to 25 alleles (*cmrDe9*) (Table 2). These two loci have also shown the lowest (*cmrDe13*) and highest (*cmrDe9*) allele numbers in other studies performed with the five markers used here (Reilly & Ward, 1999; Smith & McVeagh, 2000; Appleyard *et al.*, 2002, 2004; Shaw *et al.*, 2004; Rogers *et al.*, 2006).

Two loci (*cmrDe2* and *cmrDe9*) showed significant deviations from H-W expectations, both with deficits of heterozygotes, possibly due to the presence of null alleles or the small sample size (Table 3).

Locus *cmrDe9* has also shown significant deviation from H-W expectations in previous studies (Appleyard *et al.*, 2002, 2004; Shaw *et al.*, 2004; Rogers *et al.*, 2006), suggesting that this finding may also reflect difficulties in scoring this locus (Rogers *et al.*, 2006). In general, all null allele frequency estimators gave similar results. Only one locus showed a null allele frequency above 10% (*cmrDe2*) according to all estimators except the Brookfield-2 method, which indicated higher null allele frequencies for three loci (*cmrDe2*, *cmrDe9*, and *cmrDe30*) (Table 3). This method assumes that all individuals that fail in amplifications are null homozygotes (Brookfield, 1996). It is widely accepted that if the null allele frequency is below 10%, relatedness/parentage estimators can be used directly, without adjustment; on the other hand, loci with null allele frequencies above 50% must be excluded, due to very low power (Huang *et al.*, 2016). To avoid this distortion, and to evaluate the power for paternity/maternity exclusion analysis, two datasets were used, one with and one without genotype adjustment, with the Brookfield-2 method using Micro-Checker 2.2.3.

All genetic diversity estimates were within the ranges published in previous works on Patagonian toothfish from geographic areas other than the Chilean coast, where observed heterozygosities ranged from 0.5261 for locus *cmrDe13* (Rogers *et al.*, 2006) to 0.9480 for locus *cmrDe9* (Smith & McVeagh, 2000) (Table 2). Our findings were consistent with these previous results showing high levels of genetic diversity. In our work, all loci were strongly polymorphic with observed heterozygosities >0.6 and PIC >0.7 , indicating their potential utility in paternity exclusion analysis (Table 2).

Table 1. SSR markers and characteristics of primers used in genetic analysis of Patagonian toothfish. Underlined sequences in forward primer indicate tail to bind M13-labeled primer.

Locus	Primer sequence (5'-3')	Tm (°C)	Repeat motif	Dye	GenBank accession number
cmrDe2	F: <u>TGTA</u> AAAACGACGGCCGAGACCTCTGACAGGGTAG	66	CAA	PET	AF105071
	R: TGACAGATGTTTTCTGATTAAG	59			
cmrDe4	F: <u>TGTA</u> AAAACGACGGCCAGTATCCCAACACCAAGGCTCTATCCA	67	CAA	VIC	AF105072
	R: CCGCCTATGAGAGTCATCACGTTT	56			
cmrDe9	F: <u>TGTA</u> AAAACGACGGCCCTCTAATACTGCAGCTGTGTC	65	CA	NED	AF105073
	R: GAATCCAGAGTGTTACATAGTGA	52			
cmrDe13	F: <u>TGTA</u> AAAACGACGGCCAGAAGACAGGATAAACACACTGC	65	CA	VIC	AF105074
	R: ACCCATTTTCTTGCCCCTTTG	56			
cmrDe30	F: <u>TGTA</u> AAAACGACGGCCCTGACCTTTAACCTGCGCAAT	67	CA	VIC	AF105075
	R: ACCTCACTGATAAGGAAAGGACTG	48			

Table 2. Genetic diversity in Patagonian toothfish collected from Southern Chile. *N_a*: number of observed alleles by locus, *H_O*: observed heterozygosity, *H_E*: expected heterozygosity, *PIC*: Polymorphism information content, *PE₁*: paternity/maternity exclusion probability for a single parent, *PE₂*: paternity/maternity exclusion probability for a single parent when one parent is known, *PE_I*: identity exclusion probability for a single individual.

SSR locus	cmrDe2	cmrDe4	cmrDe9	cmrDe13	cmrDe30
N	26	32	28	34	31
<i>N_a</i>	16	12	25	8	10
Allele range size (bp)	118-180	247-283	203-267	137-161	165-185
<i>H_O</i>	0.615	0.844	0.786	0.706	0.645
<i>H_E</i>	0.912	0.869	0.966	0.788	0.804
<i>PIC</i>	0.885	0.840	0.947	0.746	0.766
H-W expectations (<i>P</i> -value)	<0.001	0.489	<0.001	0.6153	0.017
Non-adjusted genotypes					
<i>PE₁</i>	0.652	0.552	0.813	0.399	0.430
<i>PE₂</i>	0.789	0.713	0.897	0.578	0.608
<i>PE_I</i>	0.980	0.963	0.995	0.920	0.931
Adjusted genotypes					
<i>PE₁</i>	0.701	0.552	0.817	0.399	0.486
<i>PE₂</i>	0.824	0.713	0.899	0.578	0.658
<i>PE_I</i>	0.986	0.963	0.995	0.920	0.948

The global paternity exclusion probability for one putative parent using only the molecular information from progeny was high enough to perform a paternity/maternity exclusion analysis in a breeding population with five polymorphic SSR loci ($PE_1 = 0.98999$ and $PE_I = 0.99245$ with adjusted genotypes) (Araneda *et al.*, 2004). However, in a reproduction program, even if no physical tagging is performed, it is always possible to genotype all breeders. Adding genotyping information from the other parent and progeny increases the combined paternity/maternity exclusion probability. When both datasets are available, the paternity/maternity exclusion probability for a single parent can be increased to 0.99897 and 0.99927 (adjusted genotypes) when the other parent is known (PE_2). Finally, the combined identity exclusion probability was $PE_I = 0.99999998$ and $PE_I = 0.99999999$ (adjusted genotypes), indicating that these five

SSR markers show excellent potential for identifying an individual in a sample using its known genotype. While the presence of null alleles decreases the exclusion probability, a comparison of the statistics with and without adjustments reveals this reduction to be minor. In the context of a breeding population, the effect of a null allele is to mistakenly reject a true father, due to a lack of observed shared alleles with the offspring; that is, the number of real parents is underestimated. This scenario (type I error) is preferable to including a nonrelated individual as a parent (type II error) for breeding applications.

The potential application of these markers in paternity/maternity exclusion analysis must be evaluated in real families after Chilean reproduction aquaculture projects begin to produce families with viable offspring.

Table 3. Null allele frequencies for five SSR *loci* in Patagonian toothfish using six different estimators

	Van Oosterhout <i>et al.</i> (2004)	Chakraborty <i>et al.</i> (1992)	Brookfield (1996) Index 1	Brookfield (1996) Index 2	Kalinowski <i>et al.</i> (2006)	Dempster <i>et al.</i> (1977)
<i>cmrDe2</i>	0.1586	0.1847	0.1472	0.4519	0.1904	0.1402
<i>cmrDe4</i>	0.0067	0.0069	0.0063	0.1244	0.0067	0.0000
<i>cmrDe9</i>	0.0863	0.0941	0.0838	0.3265	0.0947	0.0842
<i>cmrDe13</i>	0.0471	0.0478	0.0399	0.0399	0.0498	0.0323
<i>cmrDe30</i>	0.0905	0.1018	0.0816	0.2385	0.1010	0.0814

ACKNOWLEDGEMENTS

This work was supported by FONDEF-CONICYT grant DA09i1002. We wish to thank our technician Raquel Becerra Vera for DNA extraction and general assistance.

REFERENCES

- Appleyard, S., R. Ward & R. Williams. 2002. Population structure of the Patagonian toothfish around Heard, McDonald and Macquarie Islands. *Antarctic Sci.*, 14: 364-373.
- Appleyard, S., R. Williams & R. Ward. 2004. Population genetic structure of Patagonian toothfish in the West Indian Ocean sector of the Southern Ocean. *CCAMLR Sci.*, 11: 21-32.
- Araneda, C., C. Correa, N. Lam, E. Uribe, M. Camiruaga & P. Iturra. 2004. Asignación de paternidad-maternidad y parentesco en un plantel de avestruces (*Struthio camelus*), utilizando seis *loci* microsatélites polimórficos. *Av. Produc. Anim.*, 29: 3-14.
- Brookfield, J. 1996. A simple new method for estimating null allele frequency from heterozygote deficiency. *Molec. Ecol.*, 5: 453-455.
- Chakraborty, R., M.D. Andrade, S. Daiger & B. Budowle. 1992. Apparent heterozygote deficiencies observed in DNA typing data and their implications in forensic applications. *An. Human Gen.*, 56: 45-57.
- Chapuis, M.-P. & A. Estoup. 2007. Microsatellite null alleles and estimation of population differentiation. *Mol. Biol. Evol.*, 24: 621-631.
- Dempster, A.P., N.M. Laird & D.B. Rubin. 1977. Maximum likelihood from incomplete data via the EM algorithm. *J. Roy. Statist. Soc. Ser. B*, 39(1): 1-38.
- Huang, K., K. Ritland, D.W. Dunn, X. Qi, S. Guo & B. Li. 2016. Estimating relatedness in the presence of null alleles. *Genetics*, 202: 247-260.
- Kalinowski, S.T., A.P. Wagner & M.L. Taper. 2006. ml-relate: a computer program for maximum likelihood estimation of relatedness and relationship. *Mol. Ecol. Notes*, 6: 576-579.
- Kalinowski, S.T., M.L. Taper & T.C. Marshall. 2007. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Mol. Ecol.*, 16: 1099-1106.
- Marko, P.B., H.A. Nance & K.D. Guynn. 2011. Genetic detection of mislabeled fish from a certified sustainable fishery. *Current Biol.*, 21: R621-R622.
- Reilly, A. & R.D. Ward. 1999. Microsatellite *loci* to determine population structure of the Patagonian toothfish *Dissostichus eleginoides*. *Mol. Ecol.*, 8: 1753-1754.
- Reyes, A., R. Kido & C.A. Moreno. 2012. Captura y mantención de *Dissostichus eleginoides* para conformar un plantel de reproductores. *Lat. Am. J. Aquat. Res.*, 40(4): 1066-1071.
- Rogers, A.D., S. Morley, E. Fitzcharles, K. Jarvis & M. Belchier. 2006. Genetic structure of Patagonian toothfish (*Dissostichus eleginoides*) populations on the Patagonian Shelf and Atlantic and western Indian Ocean Sectors of the Southern Ocean. *Mar. Biol.*, 149: 915-924.
- Rousset, F. 2008. GENEPOP '007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Mol. Ecol. Res.*, 8: 103-106.
- Schuelke, M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nat. Biotechnol.*, 18: 233-234.
- Shaw, P.W., A.I. Arkhipkin & H. Al-Khairulla. 2004. Genetic structuring of Patagonian toothfish populations in the Southwest Atlantic Ocean: the effect of the Antarctic Polar Front and deep-water troughs as barriers to genetic exchange. *Mol. Ecol.*, 13: 3293-3303.
- Smith, P. & M. McVeagh. 2000. Allozyme and microsatellite DNA markers of toothfish population structure in the Southern Ocean. *J. Fish Biol.*, 57: 72-83.
- Taggart, J.B., R.A. Hynes, P.A. Prodöuhl & A. Ferguson. 1992. A simplified protocol for routine total DNA isolation from salmonid fishes. *J. Fish Biol.*, 40: 963-965.
- Van Oosterhout, C., W.F. Hutchinson, D.P.M. Wills & P. Shipley. 2004. Micro-checker: software for identifying

and correcting genotyping errors in microsatellite data.
Mol. Ecol. Notes, 4: 535-538.

Received: 26 August 2015; Accepted: 4 April 2016