CLINICAL REPORTS

Complex rehabilitation of atrophic mandible with implant-supported prosthesis

Salomão Israel Monteiro Lourenço Queiroz^{1*}, Anderson Nicolly Fernandes-Costa¹, Bruno Cesar de Vasconcelos Gurgel¹, Carla Martins de Carvalho¹, Rodrigo Fromer¹, Saulo Hilton Botelho Batista¹

1. Departamento de Odontologia / Universidade Federal do Rio Grande do Norte, Brasil.

* Corresponding Author: Salomão Israel Monteiro Lourenço Queiroz

Av. Sen. Salgado Filho, 1787 - Lagoa Nova, Natal - RN, 59056-000 | Tel.: 55(84)99913-0485 | Email: salomaoisrael10@gmail.com

Work received on 19/03/2017. Approver for publication on 01/11/2017

ABSTRACT

In this study, we report the case of a patient presenting a severe atrophic mandible that was treated with short implants and stable internal fixation, without using bone grafts or biomaterials. The patient had a long history of failure of several previous consecutive treatments. Cone-beam tomography revealed a small amount of mandibular bone, suggesting a high risk of fracture of the mandible during drilling and installation of osseointegrated implants. We opted to place 5 implants (4.0 × 7.0 mm) between the mental foramina. Since the mandible presented a high risk of fracture, we inserted a titanium plate (2.3 mm) as bone reinforcement along the entire length of the mandible, including the bilateral mandibular angles. A Branemark type prosthesis was installed 5 days after the procedure. After 24 months, the patient continues to be satisfied and presents good masticatory functions without any complications.

KEYWORDS

Dental implantation, Mouth rehabilitation, Surgery, Oral.

Rev. Clin. Periodoncia Implantol. Rehabil. Oral Vol. 11(1); 47-48, 2018.

INTRODUCTION

In a normal human, the volume of cortical bone is about 3.5 times smaller than that of spongy bone, but with loss of teeth, a jaw greatly reduces this proportion as a result of vertical resorption. This cortical and spongy portion of the mandible when affected, in order to obtain greater transparency and safety⁽¹⁾.

Severe resorption of the mandible is a challenge to reestablish satisfactory masticatory function as well as harmonious functioning of the maxillofacial complex⁽²⁾. The maladaptation of a total prosthesis due to insufficient existing support causes difficulty in chewing, pain, and discomfort. Moreover, it can limit and impair the social relationships of the patients⁽³⁾. Here, we report a clinical case of a patient presenting severe mandibular resorption treated using dental implants, without mandibular bone grafts and with bone reinforcement provided using a plate and screws as a safe alternative to rehabilitation.

CASE REPORT

A 64-year-old woman without systemic diseases underwent to maxillofacial surgery and implantation at Face Defects Hospital from Sao Paulo, Brazil. She complained of masticatory problems, pain in the facial muscles, and social life limitation caused by reduced masticatory function and prosthesis maladaptation. Several previous attempts at creating an adaptable total prosthesis for her had been unsuccessful. A clinical examination, associated with cone-beam tomography, revealed a severe atrophic mandible (fig.1). Because of the imminent risk of fracture of the mandible, we treated the atrophic mandible by using 5 dental implants (4.0 × 7.0 mm, Osseotite®; Biomet-3i) and a bone reinforcement plate (2.3 mm, ThreadLock®, KLS Martin) (fig. 2, 3).

The plate was completely constructed before surgery by using a model prototype. Five days after surgery, a Branemark-type prosthesis was installed. Two years after the surgical procedure, the patient presents excellent masticatory function and is quite satisfied with the treatment (fig. 4). No postoperative complications were observed and marginal bone changes around implants were almost imperceptible with two years of follow-up.

DISCUSSION

Different surgical procedures have been described in the literature to rehabilitate patients with atrophic mandible, including onlay grafts and different types of osteotomies. The success rate of mandibular bone grafting is not high because of the high reabsorption rate of the grafts⁽⁴⁾.

Figure 1. Severe atrophic mandible: 3-D reconstruction.

Although dental implantation has been reported in atrophic mandible⁽⁵⁾, it results in serious complications, such as fracture of the mandibular bone^(6,7). Other methods used to reconstruct the mandible are osteogenic distraction, autogenous grafts, and the use of rhBMP-2. However, the outcomes of these techniques are not highly predictable and they can be quite expensive⁽⁸⁾.

Procedures of bone grafting to increase the mandible with subsequent installation of dental implants are currently the choice treatments of many researchers for reconstructing atrophic mandibles. These techniques result in survival rates of 88% to 100%⁽²⁾. The main complications reported after mandible grafting are suture dehiscence, sensory disorders, infection, and increased morbidity because 2 different surgical sites are involved⁽⁹⁾.

Another technique used to reconstruct the mandible is osteogenic distraction. The main advantage of this procedure over autogenic bone grafting is that it does not need a donor site to decrease morbidity. Another significant advantage is that osteogenic distraction allows the expansion of soft tissues⁽²⁾. However, this procedure can result in serious complications, such as decrease in the size of the previously

Figure 2. Prototype model and reconstruction plate.

Figure 3. Reconstruction plate fixed and five implants placed in mandible.

Figure 4. 24-month radiographic follow-up.

existing residual bone, infections, dehiscence of soft tissues, and loss of implants⁽¹⁰⁾.

Dental implantation is a simple surgical procedure that produces excellent results, and hence, it has become an attractive treatment⁽¹¹⁾. The main complication that could have occurred in the present case was the fracture of the atrophic mandible during implant placement. To avoid this complication, a 2.4-mm plate was installed in the mandibular bone before placing the 5 osseointegrated implants. We used a rapid prototyping model to assist in the preoperative construction of the plate as well as in surgical planning.

The virtual surgical planning of the location of the implants and their reproduction with guided surgery is an alternative during rehabilitation with dental implants⁽¹²⁾. However, the technique has some limitations, such as a structure of anatomical structures and alveolar bone with a greater risk of perforation or fenestration in cortical areas⁽¹³⁾.

The plate was placed via an intraoral approach, even though the main complications have been observed when the plate is passed bilaterally under the mental nerve bundles. Taking this into account⁽¹¹⁾, the prototype was essential for inserting the plate via a transoral approach, and this helped prevent any skin incision. After 5 days, the prosthesis was installed. We conclude that this technique leads to less surgical morbidity and low risk of infection. However, future prospective studies are needed to compare this procedure with other treatment modalities.

References

1. Blahout RM, Hienz S, Solar P, et al: Quantification of bone resorption in the interforaminal region of the atrophic mandible. *Int J Oral Maxillofac Implants* 2007;22:609-615.
2. Stellingsma C, Vissink A, Meijer HJ, Kuiper C, Raghoobar GM. Implantology and 90 the severely resorbed edentulous mandible. *Crit Rev Oral Biol Med*. 2004;15: 240-248.
3. Van Waas MA. The influence of clinical variables on patient satisfaction with complete dentures. *J Prosthet Dent*. 1990;63:307-310.
4. Bell RB, Blakey GH, White RP, Hillebrand DG, Molina A. Staged reconstruction of the severely atrophic mandible with autogenous bone graft and endosteal implants. *J Oral Maxillofac Surg* 2002;60:1135-1141.
5. Keller EE, Desjardins RP, Tolman DE, Laney WR, Van Roekel NB. Reconstruction of the severely resorbed mandibular ridge using the tissue-integrated prosthesis. *Int J Oral Maxillofac Implants* 1986;1:101-109.
6. Raghoobar GM, Stellingsma K, Batenburg RH, Vissink A. Etiology and management of mandibular fractures associated with endosteal implants in the atrophic mandible. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2000;89:553-559.
7. Marx RE, Shellenberger T, Wimsatt J, Correa P. Severely resorbed mandible: predictable reconstruction with soft tissue matrix expansion (tent pole) grafts. *J Oral Maxillofac Surg*. 2002;60:878-888.
8. Misch CM. Bone augmentation of the atrophic posterior mandible for dental implants using rhBMP-2 and titanium mesh: clinical technique and early results. *Int J Periodontics Restor Dent*. 2011;31: 581-589.
9. Tolman DE. Reconstructive procedures with endosseous implants in grafted bone: a review of the literature. *Int J Oral Maxillofac Implants* 1995;10:275-294.
10. Perdijk FB, Meijer GJ, Strijen PJ, Koole R. Complications in alveolar distraction osteogenesis of the atrophic mandible. *Int J Oral Maxillofac Surg*. 2007;36:916-921.
11. Lopes NMA, Oliveira DM, Fernandes AV, Pita Neto IC, Bezerra TP, Vasconcelos RJH. A new approach for reconstruction of a severely atrophic mandible. *J Oral Maxillofac Surg*. 2009;67:2455-2459.
12. Lemos Gulinelli Jéssica, Ferreira Edilson José, Rikio Kuabara Marcos, Borges Mattos Thiago, Borges Mattos João, Germano Evandro José et al. Accuracy of computer-guided surgery. *Rev. Clin. Periodoncia Implantol. Rehabil. Oral*. 2016, 9: 91-94.
13. Khorshid HEA, Hamed HAF, Aziz EA. Complications, risk factors, and failures of immediate functional loading of implants placed in the completely edentulous maxillae: a report of 3 consecutive cases. *Implant Dent*. 2014; 23: 125-31.