Bipartite theory of irredundant set

V. SWAMINATHAN
S. N. COLLEGE, INDIA
and

Y. B. VENKATAKRISHNAN
SASTRA UNIVERSITY, INDIA

Received : June 2010. Accepted : December 2010

Abstract

The bipartite version of irredundant set, edge-vertex irredundant set and vertex-edge irredundant set are introduced. Using the bipartite theory of graph, $IR_{ve}(G) + \gamma(G) \leq |V|$ and $\gamma_{ve}(G) + IR(G) \leq |V|$ are proved.

AMS classification : 05C69

Keywords : *Bipartite graph, X -irredundant set, Hyper Y -irredundant set, edge-vertex and vertex-edge irredundant sets.*

1. Introduction

All graphs considered here are simple and undirected. [4,5] suggests that given any problem, say P , on an arbitrary graph G , there is very likely a corresponding problem Q on a bipartite graph G^1 , such that a solution for Q provides a solution for P . The bipartite theory of graphs was introduced in [4] and a parameter called X -domination number of a bipartite graph was defined. Let $G = (X, Y, E)$ be a bipartite graph with $|X| = p$ and $|Y| = q$. Two vertices u and v in X are X -adjacent if they have a common adjacent vertex $y \in Y$. Let $y \in X$ and $\Delta_Y = \max\{|N_Y(u)| : y \in X\}$ where the X -neighbor set $N_Y(u)$ is defined as $N_Y(u) = \{v \in X : u \text{ and } v \text{ are } X\text{-adjacent}\}$.

A subset $X \subseteq X$ is an X -dominating set [4] if every $x \in X - D$ is X -adjacent to some vertex in D . The minimum cardinality of a X -dominating set is called X -domination number and is denoted by $\gamma_X(G)$.

We say a vertex $x \in X$ hyper Y -dominates $y \in Y$ if $y \in N(x)$ or $y \in N(N_Y(x))$. A subset $S \subseteq X$ is a hyper Y -dominating set [6] if every $y \in Y$ is hyper Y -dominated by a vertex of S . The minimum cardinality of a hyper Y -dominating set is called hyper Y -domination number and is denoted by $\gamma_{hY}(G)$.

Given an arbitrary graph $G = (V, E)$, a vertex $u \in V(G)$ ve-dominates an edge $vw \in E(G)$ if (a) $u = v$ or $u = w$ (u incident to vw) or (b) uv or uw is an edge in G . A subset $S \subseteq V(G)$ is a vertex-edge dominating set [3] if for all edges $e \in E(G)$, there exists a vertex $v \in S$ such that v dominates e . The minimum cardinality of a ve-dominating set of G is called the vertex-edge domination number and is denoted as $\gamma_{ve}(G)$.

An edge $e = uv \in E(G)$ ev-dominates a vertex $w \in V(G)$ if (i) $u = w$ or $v = w$ (w is incident to e) or (ii) uw or vw is an edge in G . (w is adjacent to u or v). A set $S \subseteq E(G)$ is an edge-vertex dominating set [3] if for all vertices $v \in V(G)$, there exists an edge $e \in S$ such that e dominates v . The minimum cardinality of a ev-dominating set of G is called the edge-vertex domination number and is denoted as $\gamma_{ev}(G)$.

Observation: 1. Let G be an arbitrary graph. A vertex $u \in V(G)$ ve-dominates the edge $e \in E(G)$ if and only if the edge e ev-dominates the vertex $u \in V(G)$.

2. Bipartite Construction

The bipartite graph $VE(G)$ constructed from an arbitrary graph $G = (V, E)$ is defined as in [4]. $VE(G) = (V, E, F)$ is defined by the edges $F = \{(u, e) : e = (u, v) \in E\}$. $VE(G) \cong S(G)$, where $S(G)$ denotes the subdivision graph of G .

The bipartite graph $EV(G)$ [4] constructed from an arbitrary graph $G = (V, E)$ is defined as $EV(G) = (E, V, J)$ where $J = \{(e, u)(e, v) : e = (u, v) \in E\}$.

A set $S \subseteq V$ of vertices in a graph $G = (V, E)$ is called a dominating set [2] if every $v \in V$ is either an element of S or is adjacent to an element of S . The minimum cardinality of a dominating set of a graph G is called the domination number and is denoted by $\gamma(G)$.

A set $F \subseteq E(G)$ of edges in a graph $G = (V, E)$ is called an edge dominating set [2] if every $e \in E(G)$ is either an element of F or is adjacent to an element of $E - F$. The minimum cardinality of an edge dominating set of a graph G is called the edge domination number and is denoted by $\gamma_1(G)$.

Theorem:2.1 [4] For any graph G ,

- (a) $\gamma_X(VE(G)) = \gamma(G)$
- (b) $\gamma_X(EV(G)) = \gamma_1(G)$.

Theorem:2.2 [6] For any graph G ,

- (a) $\gamma_{hY}(VE(G)) = \gamma_{ve}(G)$
- (b) $\gamma_{hY}(EV(G)) = \gamma_{ev}(G)$.

3. Irredundant sets

3.1. Vertex-edge irredundant set

A vertex $v \in S \subseteq V(G)$ has a private edge $e = uv \in E(G)$ (with respect to a set S), if: 1. v is incident to e or v is adjacent to either u or w , and 2. for every vertices $x \in S - \{v\}$, x is not incident to e and x is not adjacent to either u or w .

A set S is a vertex-edge irredundant set [3] (simply a ve-irredundant set) if every vertex $v \in S$ has a private edge. The vertex-edge irredundance of a graph G is the cardinality of a maximal ve-irredundant set with minimum number of vertices and is denoted by $ir_{ve}(G)$. The upper vertex-edge irredundance number of a graph G is the cardinality of a maximum

ve-irredundant set of vertices and is denoted by $IR_{ve}(G)$.

Theorem: 3.1.1 [3] Every minimal ve-dominating set is a maximal ve-irredundant set.

3.2. Edge-vertex irredundant set

An edge $e = uv \in F \subseteq E(G)$ has a private vertex $w \in V(G)$ (with respect to a set F), if: 1. e is incident to w , and 2. for all edges $f = xy \in F - \{e\}$, f is not incident to w and neither x nor y is adjacent to w .

A set F is an edge-vertex irredundant set [3] (simply a ev-irredundant set) if every edge $e \in F$ has a private vertex. The edge-vertex irredundance of a graph G is the cardinality of a maximal ev-irredundant set with minimum number of vertices and is denoted by $ir_{ev}(G)$. The upper edge-vertex irredundance number of a graph G is the cardinality of a maximum ev-irredundant set of vertices and is denoted by $IR_{ev}(G)$.

Theorem 3.2.1:[3] Every minimal ev-dominating set of G is a maximal ev-irredundant set.

3.3. Hyper Y - Irredundant set

Let $G = (X, Y, E)$ be a bipartite graph. Let $S \subseteq X$. A vertex $x \in S$ has a private hyper Y -neighbor $y \in Y$ if 1. x is adjacent to y or $y \in N(N_Y(x))$ and 2. for all vertices $x_1 \in S - \{x\}$, x_1 is not adjacent to y and $y \notin N(N_Y(x_1))$.

A set S is hyper Y -irredundant set if every $v \in S$ has a private hyper Y -neighbor. The hyper Y -irredundance number of a graph G is the minimum cardinality of a maximal hyper Y -irredundant set of vertices and is denoted by $ir_{hY}(G)$. The upper hyper Y -irredundance number of a graph G is the maximum cardinality of a maximal hyper Y -irredundant set of vertices and is denoted by $IR_{hY}(G)$.

Theorem: 3.3.1 A hyper Y -dominating set S is a minimal hyper Y -dominating set if and only if it is hyper Y -dominating set and hyper Y -irredundant set.

Proof: Let S be a hyper Y -dominating set. Then S is a minimal hyper Y -dominating set if and only if $\forall u \in S, \exists y \in Y$ which is not hyper Y -dominated by $S - \{u\}$. Equivalently, S is a minimal hyper Y -dominating set if and only if $\forall u \in S, u$ has atleast one private hyper Y -neighbour. Thus S is minimal hyper Y -dominating set if and only if it is hyper Y -irredundant set.

Conversely, let S be both hyper Y -dominating and hyper Y -irredundant.

Claim: S is a minimal hyper Y -dominating set.

If S is not minimal hyper Y -dominating set, there exists $v \in S$ for which $S - \{v\}$ is hyper Y -dominating. Since S is hyper Y -irredundant, v has a private hyper Y -neighbor of u . By definition u is not hyper Y -adjacent to any vertex in $S - \{v\}$. That is, $S - \{v\}$ is not hyper Y -dominating set, a contradiction. Hence, S is a minimal hyper Y -dominating set.

Theorem: 3.3.2 Every minimal hyper Y -dominating set is a maximal hyper Y -irredundant set.

Proof: Every minimal hyper Y -dominating set S is hyper Y -irredundant set.

Claim: S is a maximal hyper Y -irredundant set.

Suppose S is not maximal hyper Y -irredundant set. Then there exists a vertex $u \in X - S$ for which $S \cup \{u\}$ is hyper Y -irredundant. There exists atleast one vertex $y \in Y$ which is a private hyper Y -neighbor of u with respect to $S \cup \{u\}$. That is no vertex in S is hyper Y -adjacent to y . Hence, S is not a hyper Y -dominating set, a contradiction. Hence, S is a maximal hyper Y -irredundant set.

Theorem:3.3.3 For any graph G ,

$$(a) \text{ } ir_{hY}(VE(G)) = ir_{ve}(G)$$

$$(b) \text{ } ir_{hY}(EV(G)) = ir_{ev}(G).$$

Proof: Let S be a ir_{hY} -set of $VE(G) = (X, Y, E)$. Every $x \in S$ has a private hyper Y -neighbor $y \in Y$. x is adjacent to y or $y \in N(N_Y(x))$ and for all vertices $x_1 \in S - \{x\}$, x_1 is not adjacent to y and $y \notin N(N_Y(x_1))$. In graph G , $x \in S \subseteq V$ is incident with $y \in E$ or x is adjacent to either u or v where $y = uv$ and for every $x_1 \in S - \{x\}$, $y \in E$ is not incident with

x_1 and x_1 is not adjacent to either u or v . S is a vertex edge irredundant set.

$$ir_{ev}(G) \leq |S| = ir_{hY}(VE(G)).$$

Let U be a ir_{ve} -set of G . Every vertex $v \in S$ has a private edge $e = uv$ with respect to U . Equivalently, v is incident with e or v is adjacent to either u or w and for every $x \in U - \{v\}$, x is not incident with e and x is not adjacent to either u or w . In $VE(G)$, every $v \in S$ has private hyper Y -neighbor e . Therefore, $U \subseteq X$ is a hyper Y -irredundant set of $VE(G)$. Hence, $ir_{hY}(VE(G)) \leq |U| = ir_{ve}(G)$.

Similarly (b) can be proved.

3.4. X-Irredundant set

Let $G = (X, Y, E)$ be a bipartite graph. Let $S \subseteq X$. Let $u \in S$. A vertex v is a private X -neighbor of u with respect to S if u is the only point of S , X -adjacent to v .

A set S is X -irredundant set if every $u \in S$ has a private X -neighbor. The X -irredundance number of a graph G is the cardinality of a maximal X -irredundant set of vertices with minimum cardinality and is denoted by $ir_X(G)$. The upper X -irredundance number of a graph G is the cardinality of a X -irredundant set of vertices with maximum cardinality and is denoted by $IR_X(G)$.

Theorem:3.4.1 A X -dominating set S is a minimal X -dominating set if and only if it is X -dominating and X -irredundant.

Proof: Let S be a X -dominating set. Then S is a minimal X -dominating set if and only if for every $u \in S$ there exists $v \in X - (S - \{u\})$ which is not X -dominated by $S - \{u\}$. Equivalently, S is a minimal X -dominating set if and only if $\forall u \in S$, u has atleast one private X -neighbor with respect to S . Thus S is minimal X -dominating set if and only if it is X -irredundant.

Conversely, Let S is both X -dominating and X -irredundant.

Claim: S is a minimal X -dominating set.

If S is not a minimal X -dominating set, then there exists $v \in S$ for which $S - \{v\}$ is X -dominating. Since S is X -irredundant, v has a private X -neighbor of with respect to S say u (u may be equal to v). By definition, u is not X -adjacent to any vertex in $S - \{v\}$. Therefore, $S - \{v\}$ is not a X -dominating set, a contradiction. Hence, S is a minimal X -dominating

set.

Theorem:3.4.2 Every minimal X -dominating set is a maximal X -irredundant set.

Proof: Every minimal X -dominating set S is X -irredundant set.

Claim: S is a maximal X -irredundant set.

Suppose S is not a maximal X -irredundant set. Then there exists a vertex $u \in X - S$ for which $S \cup \{u\}$ is X -irredundant. Therefore, there exists atleast one vertex x which is a private X -neighbor of u with respect to $S \cup \{u\}$. Hence, no vertex in S is X -adjacent to x . Thus S is not X -dominating set, a contradiction. Hence, S is maximal X -irredundant set.

A vertex v is a private neighbor of a vertex u in a set $S \subseteq V(G)$ with respect to S if $N[v] \cap S = \{u\}$. The private neighbor set of $u \in S$ with respect to S is defined as $pn[u, S] = \{v : N[v] \cap S = \{u\}\}$. A set S is called irredundant set [2] if for every vertex $u \in S$, $pn[u, S] \neq \phi$. The irredundance number of agraph G is the cardinality of a maximal irredundant set with minimum number of vertices and is denoted by $ir(G)$. The upper irredundance number of a graph G is the cardinality of a maximum irredundant set of vertices and is denoted by $IR(G)$.

Theorem:3.4.3 For any graph G ,

- (a) $ir_X(VE(G)) = ir(G)$
- (b) $ir_X(EV(G)) = ir^1(G)$

Proof: Let S be a ir_X set of $VE(G) = (X, Y, E^1)$. Every v has a private X -neighbor u . Equivalently, v is X -adjacent to u and no other vertex in S is X -adjacent to u . In G , $v \in S$ is the only vertex adjacent to u and no other vertex in S is adjacent to u . Therefore, S is an irredundant set of G .

$$ir(G) \leq |S| = ir_X(VE(G)).$$

Let U be an ir - set of G . For every vertex $v \in U$, $pn[v, U] \neq \phi$. Every vertex $v \in U$ has at least one private neighbor with respect to u . In $VE(G)$, that is every vertex $v \in U$ has at least one private X -neighbor. Therefore, U is an X -irredundant set. Hence, $ir_X(VE(G)) \leq |U| = ir(G)$. Hence, $ir_X(VE(G)) = ir(G)$.

(b) Let S be an ir_X set of $EV(G) = (X, Y, E^1)$. Every e has a private X -neighbor f . Equivalently, e is X -adjacent to f and no other vertex in S is X -adjacent to f . In G , $e \in S$ is the only edge adjacent to f and no other edge in S is adjacent to f . Therefore, S is an edge irredundant set of G . Hence, $ir^1(G) \leq |S| = ir_X(EV(G))$.

Let U be a ir^1 -set of G . For every edge $e \in U$, $pn[e, U] \neq \phi$. Hence, every edge $e \in U$ has at least one private neighbor. That is, in $EV(G)$, every vertex $e \in U$ has at least one private X -neighbor. Therefore, U is an X -irredundant set in $EV(G)$. Thus, $ir_X(EV(G)) \leq |U| = ir^1(G)$. Hence, $ir_X(EV(G)) = ir^1(G)$.

4. Main Result

For any graph G , $IR_{ve}(G) + \gamma(G) \leq |V|$ and $\gamma_{ve}(G) + IR(G) \leq |V|$ are proved using bipartite theory of graphs, which are open problem in [3].

Theorem:4.1 Let $G = (X, Y, E)$ be a bipartite graph with $N_Y(x) \neq \phi$ for every $x \in X$. Then $IR_{hY}(G) + \gamma_X(G) \leq |X|$.

Proof: Let S be a IR_{hY} set of G . Then, S is a maximal hyper Y -irredundant set. Therefore, S is a hyper Y -irredundant set. That is every $x \in S$ has a private hyper Y -neighbor $y \in Y$. Then x is adjacent to y or $y \in N(N_Y(x))$ and for all vertices $x_1 \in S - \{x\}$, x_1 is not adjacent to y and $y \notin N(N_Y(x))$.

Case(i): x is adjacent with y .

Since $N_Y(v) \neq \phi$, x has X -neighbours. Let z be any X -neighbour of x . Suppose $z \in S$. Then z is not adjacent to y and $y \notin N(N_Y(z))$. But $y \in N(N_Y(x))$, since x is a X -neighbour of z , a contradiction. Therefore, any X -neighbour of x is in $X - S$.

Case(ii): $y \in N(N_Y(x))$.

Vertices in $N(y)$ are in $X - S$. Then $N(y) \subseteq X - S$. Other wise, we get a contradiction to $y \in Y$ is a private hyper Y -neighbor of $x \in S$. Hence, for every $x \in S$ there exists $x_1 \in X - S$ such that x and x_1 are X -adjacent. That is, $X - S$ is a X -dominating set. Therefore, $\gamma_X(G) \leq |X - S| = |X| - IR_{hY}(G)$. Hence, $IR_{hY}(G) + \gamma_X(G) \leq |X|$.

Corollary: 4.2 For any graph G ,

$$(a) IR_{ve}(G) + \gamma(G) \leq |V|$$

$$(b) IR_{ev}(G) + \gamma_1(G) \leq |E|.$$

Theorem:4.3 Let $G = (X, Y, E)$ be a bipartite graph with $N_Y(x) \neq \phi$ for every $x \in X$ then $IR_X(G) + \gamma_{hY}(G) \leq |X|$.

Proof: Let S be a IR_X set of G . Every element $x \in S$ has a private X -neighbor. Consider the set $X - S$. Since $X - S$ is a X -dominating set elements of Y are either adjacent to $X - S$ or adjacent to vertices which are X -adjacent to elements of $X - S$. Therefore, $X - S$ is a hyper Y -dominating set. Therefore, $\gamma_{hY} \leq |X - S| = |X| - IR_X$. Hence, $IR_X + \gamma_{hY} \leq |X|$.

Corollary: 4.4 For any graph G ,

$$(a) \gamma_{ve}(G) + IR(G) \leq |V|$$

$$(b) \gamma_{ev}(G) + IR^1(G) \leq |E|.$$

Acknowledgement: We are thankful to the anonymous referee for helpful suggestions, which led to substantial improvement in the paper.

References

- [1] Bondy J. A., Murthy U. S. R., *Graph theory with applications*, London Macmillan (1976).
- [2] Haynes T. W., Hedetniemi. S. T. and Slater P. J., *Fundamentals of Domination in graphs*, Marcel Dekker, New York, (1998).
- [3] Jason Robert Lewis, *Vertex-edge and edge-vertex parameters in graphs*, (Ph. D Thesis), Clemson University, August 2007.
- [4] Stephen Hedetniemi, Renu Laskar, *A Bipartite theory of graphs I*, Congressus Numerantium, Volume **55**; pp. 5–14, December 1986.
- [5] Stephen Hedetniemi, Renu Laskar, *A Bipartite theory of graphs II*, Congressus Numerantium, Volume **64**; pp. 137-146, November 1988.

- [6] Swaminathan V. and Venkatakrishnan Y. B., *Hyper Y-domination in Bipartite graphs*, International Mathematical Forum, Volume **4**, No. **20**, pp. 953-958, (2009).

V. Swaminathan^a

Research Coordinator,
Ramanujan Research Centre,
S. N. College,
Madurai,
India
e-mail : sulanesri@yahoo.com

and

Y. B. Venkatakrishnan^b

Department of Mathematics,
SASTRA University,
Tanjore,
India
e-mail : venkatakrish2@maths.sastra.edu