An extension of Sheffer polynomials

A. K. SHUKLA

S. V. NATIONAL INSTITUTE OF TECHNOLOGY, INDIA

S. J. RAPELI

S. V. NATIONAL INSTITUTE OF TECHNOLOGY, INDIA

Received : February 2011. Accepted : September 2011

Abstract

Sheffer [Some properties of polynomial sets of type zero, Duke Math. J. 5 (1939), pp.590-622] studied polynomial sets zero type and many authors investigated various properties and its applications. In the sequel to the study of Sheffer Polynomials, an attempt is made to generalize the Sheffer polynomials by using partial differential operator.

Keywords : *Appell sets, Differential operator, Sheffer polynomials, generalized Sheffer polynomials.*

AMS Subject Classification (2010): *33C65, 33F99, 44A45, 46G25.*

1. Introduction

Sheffer [8] wrote remarkable paper on some properties of polynomials of type zero, Huff and Rainville [4] obtained generating function of Sheffer Polynomial and Rainville [7] reported this work in his book. Al Salam and Verma [1] studied the generalized Sheffer polynomials and Bahman Alidad [2] studied Sheffer polynomials in two variables. Recently Galiffa Joseph [3] worked on Sheffer B-type 1 orthogonal polynomial sequences in his thesis.

A polynomial set $\{P_n\}_{n \geq 0}$ is said to be of Sheffer A-type zero [5] if and only if it has a generating function of the form

$$A(t) \exp(xG(t)) = \sum_{n=0}^{\infty} P_n(x)t^n,$$

where $A(t)$ and $G(t)$ are two formal power series

$$A(t) = \sum_{k=0}^{\infty} a_k t^k, \quad a_0 \neq 0, \quad G(t) = \sum_{k=0}^{\infty} g_k t^{k+1}, \quad g_0 \neq 0.$$

This has the property: $J(D)P_0(x) = 0$ and $JP_n(x) = P_{n-1}(x)$, $n \geq 1$; where $J(D)$ is defined by

$$J = J(D) = \sum_{k=0}^{\infty} a_k D^{k+1}, \quad a_0 \neq 0 \quad \text{and} \quad D = \frac{d}{dx}.$$

Bahman Alidad [2] studied polynomials in two variables of Sheffer A-type zero and also proved some related theorems. Sheffer [8] and Steffensen [9] generalized Appell sets by considering a linear differential operator of infinite order with constant coefficients

$$L(D)P_n(x) = P_{n-1}(x), \quad \text{where} \quad L(D) = \sum_{k=0}^{\infty} c_k D^{k+1}, \quad c_0 \neq 0.$$

Osegove [6] has also generalized Appell sets and studied polynomial sets which has the property $D^r P_n(x) = P_{n-r}(x)$, $(n = r, r+1, \dots)$, where r is a fixed positive number.

Al Salam and Verma [1] generalized Sheffer- Steffensen polynomials and defined as

$$J(D)P_n(x) = P_{n-r}(x), \quad n = r, r+1, \dots, \quad \text{where} \quad J(D) = \sum_{k=0}^{\infty} a_k D^{k+r}, \quad a_0 \neq 0$$

and $a_k (k \geq 0)$ is independent of x .

In sequel to the study, we consider a class of polynomials that contain both the Sheffer-Steffensen sets and the Osego sets of polynomials in two variables.

Let $\{P_n(x, y)\}$ be a simple set polynomials in two variables x and y , i.e. a sequence of polynomials $P_0(x, y), P_1(x, y), P_2(x, y), \dots$ for which the degree of $P_n(x, y)$ is exactly n .

A polynomial set in two variables is said to be Appell set if $P'_n(x, y) = P_{n-1}(x, y)$, for all $n \geq 1$. A well known characteristic property of Appell set is the existence of a (formal) power series $A(t) = \sum_{n=0}^{\infty} a_n t^n$, $a_0 \neq 0$ such that

$$(1.1) \quad \sum_{n=0}^{\infty} P_n(x, y) t^n = A(t) \exp(xt) \exp(yt)$$

In generalized Appell sets, consider a linear differential operator of infinite order with constant coefficients

$$(1.2) \quad L(D) = \sum_{k=0}^{\infty} C_k D^{k+1}, \quad C_0 \neq 0,$$

as a generalization of the differential operator $D = \frac{\partial}{\partial x} + \frac{\partial}{\partial y}$.

The polynomial set which satisfy the following property

$$(1.3) \quad L(D)P_n(x, y) = P_{n-1}(x, y),$$

Bahman Alidad [2] proved that polynomial sets have the property as Equation (1.3) if and only if polynomial sets have the generating function in following form

$$(1.4) \quad A(t) \exp(xG(t)) \exp(yH(t)) = \sum_{n=0}^{\infty} P_n(x, y) t^n$$

where

$$G(t) = \sum_{n=0}^{\infty} g_n t^{n+1}, \quad g_0 \neq 0,$$

$$H(t) = \sum_{n=0}^{\infty} h_n t^{n+1}, \quad h_0 \neq 0,$$

$$A(t) = \sum_{n=0}^{\infty} a_n t^n, \quad a_0 \neq 0.$$

The polynomial sets in two variables have the property as given below

$$(1.5) \quad D^r P_n(x, y) = P_{n-r}(x, y) \quad (n = r, r+1, \dots),$$

where r is a (fixed) positive integer and $D = \frac{\partial}{\partial x} + \frac{\partial}{\partial y}$.

2. The generalized Sheffer-Steffensen polynomials

Let r be a fixed positive integer and we consider the differential operator (formal) of infinite order

$$(1.6) \quad J(D) = \sum_{k=0}^{\infty} a_k D^{k+r}, \quad a_0 \neq 0, \quad D = \frac{\partial}{\partial x} + \frac{\partial}{\partial y},$$

where $a_k (k \geq 0)$ is independent of x . We shall say that the polynomial set $\{P_n(x, y)\}$ is in $S^{(r)}$ if there is an operator $J(D)$ of the form Equation (1.6) such that

$$(1.7) \quad J(D)P_n(x, y) = P_{n-r}(x, y) \quad (n = r, r+1, r+2, \dots)$$

Obviously $S^{(r)}$ contains the Osego class $A^{(r)}$ of polynomial sets which satisfy Equation (1.5). We clearly see that $S^{(1)}$ is the class of Sheffer A-type zero polynomials. It is also clear that $S^{(r)}$ may be partitioned into equivalence classes $\{S_J^{(r)}\}$ where $S_J^{(r)}$ is the class of all polynomial sets with the property that

$$J(D)P_n(x, y) = P_{n-r}(x, y),$$

for a fixed operator J .

If $\{p_n(x, y)\} \in S_J^{(r)}, \{q_n(x, y)\} \in S_J^{(r)}$ then $\{\alpha p_n(x, y) + \beta q_n(x, y)\} \in S_J^{(r)}$,

where α, β are constants or are periodic in n of period r .

Theorem 2.1: Let $\{P_n(x, y)\}$ be a polynomial set in two variables. In order to that $\{P_n(x, y)\}$ be in $S^{(r)}$, it is necessary and sufficient that there exist (formal) power series

$$(1.8) \quad G(t) = \sum_{i=1}^{\infty} g_i t^i, \quad g_1 \neq 0,$$

$$(1.9) \quad H(t) = \sum_{i=1}^{\infty} h_i t^i, \quad h_1 \neq 0,$$

$$(1.10) \quad A_s(t) = \sum_{i=0}^{\infty} \alpha_i^{(s)} t^i, \quad (\text{not all } \alpha_0^{(s)} \text{ are zeros})$$

such that

$$(1.11) \quad \sum_{i=1}^r A_i(t) \exp((xG(\varepsilon_i(t))) \exp((yH(\varepsilon_i(t))) = \sum_{n=0}^{\infty} P_n(x, y) t^n$$

Proof: If $F(t) = \sum_{k=0}^{\infty} \alpha_k t^k$, $\alpha_0 \neq 0$, is a formal power series then we can find another formal power series $f(t) = \sum_{k=0}^{\infty} \beta_k t^k$, $\beta_0 \neq 0$ such that

$$(1.12) \quad F(t) = (f(t))^k, \quad k = 1, 2, \dots$$

Now let the polynomial sets $\{P_n(x, y)\} \in S^{(r)}$ and suppose that the differential operator corresponding to $\{P_n(x, y)\}$ is

$$J(D) = \sum_{k=0}^{\infty} a_k D^{k+r}, \quad a_0 \neq 0, \quad D = \frac{\partial}{\partial x} + \frac{\partial}{\partial y}.$$

We write $J(t) = t^r F(t)$ and using Equation (1.12)

$$J(t) = \{t f(t)\}^r = \{L(t)\}^r,$$

therefore, that $L(D)$ is a differential operator of the form Equation (1.2).

Let

$$F(x, y, t) = \sum_{n=0}^{\infty} P_n(x, y) t^n$$

Now

$$\begin{aligned} J(D)F(x, y, t) &= \sum_{n=0}^{\infty} J(D)P_n(x, y) t^n \\ &= \sum_{n=r}^{\infty} P_{n-r}(x, y) t^n \\ &= \sum_{n=0}^{\infty} P_n(x, y) t^{n+r} \end{aligned}$$

$$\begin{aligned}
&= t^r \sum_{n=0}^{\infty} P_n(x, y) t^n \\
J(D)F(x, y, t) &= t^r F(x, y, t) \\
\{J(D) - t^r\}F(x, y, t) &= 0
\end{aligned}$$

Consequently if $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r$ are the r roots of unity, we get that

$$\{L(D) - \varepsilon_i t\}F(x, y, t) = 0 \quad (i = 1, 2, \dots, r).$$

Hence

$$(1.13) \quad F(x, y, t) = \sum_{i=1}^r A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)),$$

where $L(H(t)) = t$.

Conversely if $F(x, y, t)$ has the form as Equation (1.13), then it follows that

$$\begin{aligned}
J(D)F(x, y, t) &= t^r F(x, y, t) \\
J(D) \sum_{n=0}^{\infty} P_n(x, y) t^n &= t^r \sum_{n=0}^{\infty} P_n(x, y) t^n \\
\sum_{n=0}^{\infty} J(D)P_n(x, y) t^n &= \sum_{n=0}^{\infty} P_n(x, y) t^{n+r} \\
\sum_{n=r}^{\infty} J(D)P_n(x, y) t^n &= \sum_{n=r}^{\infty} P_{n-r}(x, y) t^n
\end{aligned}$$

So that

$$J(D)P_n(x, y) = P_{n-r}(x, y).$$

This completes the proof of Theorem 2.1.

Theorem 2.2: Let two polynomial sequences $P_n(x, y)$ and $Q_n(x, y)$ are in $S^{(r)}$ and there is an arbitrary sequence of numbers $\{a_n\}$, $a_0 \neq 0$, if and only if

$$(1.14) \quad Q_n(x, y) = \sum_{k=0}^n a_{n-k} P_k(x, y)$$

Proof: Consider $\{P_n(x, y)\}, \{Q_n(x, y)\} \in S^{(r)}$.

Therefore $P_n(x, y)$ and $Q_n(x, y)$ belong to the same operator $J(D)$ as in form of Equation (1.6) such that

$$J(D)P_n(x, y) = P_{n-r}(x, y) \quad \text{and} \quad J(D)Q_n(x, y) = Q_{n-r}(x, y),$$

where $(n = r, r + 1, r + 2, \dots)$.

Since $P_n(x, y)$ and $Q_n(x, y)$ are simple set of polynomials [5] in two variables,

$$Q_n(x, y) = \sum_{k=0}^n a_{n,k} P_k(x, y)$$

Thus

$$\begin{aligned} \sum_{k=0}^{n-r} a_{n-r,k} P_k(x, y) &= Q_{n-r}(x, y) = JQ_n(x, y) \\ &= \sum_{k=r}^n a_{n,k} J P_k(x, y) \\ &= \sum_{k=r}^n a_{n,k} P_{k-r}(x, y) \\ &= \sum_{k=0}^{n-r} a_{n,k+r} P_k(x, y) \end{aligned}$$

Hence $a_{n-r,k} = a_{n,k+r}$ or $a_{n,k+r} = a_{n-k-r,0}$, yields Equation (1.14).

Conversely: Assume Equation (1.14) to hold. There exists an operator J to which $P_n(x, y)$ belongs. It follows from Equation (1.14) that $Q_n(x, y)$ also belongs to same operator J . This proves Theorem 2.2.

Theorem 2.3: Let $P_n(x, y)$ be of Sheffer A-type zero, $P_n(x, y)$ belong to the operator $J(D)$ and have the generating function in equation (1.11) of Theorem 2.1. There exist sequences $\alpha_k^{(s)}$, μ_k and η_k , independent of x, y and n , such that for all $n \geq 1$,

$$(1.15) \quad nP_n(x, y) = \sum_{i=1}^r S_{i,n}(x, y) \psi(\varepsilon_i t),$$

where $S_{i,n}(x, y) = \alpha_{n-1}^{(i)} + x\varepsilon_i^n \mu_{n-1} + y\varepsilon_i^n \eta_{n-1}$ and $\psi(\varepsilon_i t) = A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t))$ and $\mu_k = (k+1)g_k$ in terms of g_k of Equation (1.8) and $\eta_k = (k+1)h_k$, in terms of h_k of Equation (1.9).

Proof: Consider the series

$$\sum_{n=0}^{\infty} P_n(x, y) t^n = \sum_{i=1}^r A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t))$$

and writing this in the following form

$$t \frac{\partial}{\partial t} \left(\sum_{n=0}^{\infty} P_n(x, y) t^n \right) = t \frac{\partial}{\partial t} \left(\sum_{i=1}^r A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)) \right)$$

or

$$\sum_{n=0}^{\infty} n P_n(x, y) t^n = t \left[\sum_{i=1}^r \left\{ A_i'(t) + x \varepsilon_i G'(\varepsilon_i t) A_i(t) + y \varepsilon_i H'(\varepsilon_i t) A_i(t) \right\} \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)) \right]$$

On simplifying above equation, yields

$$\begin{aligned} \sum_{n=0}^{\infty} n P_n(x, y) t^n &= t \left[\sum_{i=1}^r \left\{ \frac{A_i'(t)}{A_i(t)} + x \varepsilon_i G'(\varepsilon_i t) + y \varepsilon_i H'(\varepsilon_i t) \right\} A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)) \right] \\ &= t \left[\sum_{i=1}^r \left\{ \sum_{n=0}^{\infty} \alpha_n^{(i)} t^n + x \varepsilon_i \sum_{n=0}^{\infty} \mu_n \varepsilon_i^n t^n + y \varepsilon_i \sum_{n=0}^{\infty} \eta_n \varepsilon_i^n t^n \right\} A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)) \right] \\ &= t \left[\sum_{i=1}^r \left\{ \sum_{n=0}^{\infty} \left(\alpha_n^{(i)} + x \varepsilon_i^{n+1} \mu_n + y \varepsilon_i^{n+1} \eta_n \right) t^n \right\} A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)) \right] \\ &= \sum_{n=0}^{\infty} \sum_{i=1}^r \left(\alpha_n^{(i)} + x \varepsilon_i^{n+1} \mu_n + y \varepsilon_i^{n+1} \eta_n \right) A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)) t^{n+1} \\ &= \sum_{n=1}^{\infty} \sum_{i=1}^r \left(\alpha_{n-1}^{(i)} + x \varepsilon_i^n \mu_{n-1} + y \varepsilon_i^n \eta_{n-1} \right) A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)) t^n \end{aligned}$$

So that

$$\sum_{n=1}^{\infty} n P_n(x, y) t^n = \sum_{n=1}^{\infty} \sum_{i=1}^r S_{i,n}(x, y) \psi(\varepsilon_i t) t^n$$

Hence

$$n P_n(x, y) = \sum_{i=1}^r S_{i,n}(x, y) \psi(\varepsilon_i t),$$

where

$$S_{i,n}(x, y) = \alpha_{n-1}^{(i)} + x \varepsilon_i^n \mu_{n-1} + y \varepsilon_i^n \eta_{n-1}$$

and

$$\psi(\varepsilon_i t) = A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)).$$

This gives the statement.

Theorem 2.4: A necessary and sufficient condition that $P_n(x, y)$ be of Sheffer A-type zero, there exists sequence g_k and h_k , independent of x, y and n , such that

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right) P_n(x, y) = \sum_{i=1}^r (\varepsilon_i^n g_{n-1} + \varepsilon_i^n h_{n-1}) \psi(\varepsilon_i t)$$

Proof:

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right) \left(\sum_{n=0}^{\infty} P_n(x, y) t^n\right) = \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right) \left[\frac{\sum_{i=1}^r A_i(t) \exp(xG(\varepsilon_i t))}{\exp(yH(\varepsilon_i t))} \right]$$

$$\sum_{n=0}^{\infty} \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right) P_n(x, y) t^n = \sum_{i=1}^r \{G(\varepsilon_i t) + H(\varepsilon_i t)\} A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t))$$

$$= \sum_{i=1}^r \sum_{n=0}^{\infty} (\varepsilon_i^{n+1} g_n + \varepsilon_i^{n+1} h_n) t^{n+1} \psi(\varepsilon_i t)$$

$$= \sum_{n=1}^{\infty} \sum_{i=1}^r (\varepsilon_i^n g_{n-1} + \varepsilon_i^n h_{n-1}) \psi(\varepsilon_i t) t^n$$

Thus

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y}\right) P_n(x, y) = \sum_{i=1}^r (\varepsilon_i^n g_{n-1} + \varepsilon_i^n h_{n-1}) \psi(\varepsilon_i t),$$

where

$$\psi(\varepsilon_i t) = A_i(t) \exp(xG(\varepsilon_i t)) \exp(yH(\varepsilon_i t)).$$

Hence, we get theorem 2.4.

References

- [1] W. A. Al Salam and A. Verma, Generalized Sheffer Polynomials, Duke Math. J. 37, pp. 361-365, (1970).
- [2] B. Alidad, On some problems of special functions and structural matrix analysis, Ph.D. diss., Aligarh Muslim University, (2008).
- [3] Joseph D. Galiffa, The Sheffer B-type 1 orthogonal polynomial sequences, Ph.D. diss., University of Central Florida, (2009).
- [4] William N. Huff and E. D. Rainville, On the Sheffer A-type of polynomials generated by $\phi(t)f(xt)$, Proc. Amer. Math. Soc. 3, pp. 296-299 (1952).
- [5] E. B. Mc Bride, Obtaining Generating Functions, Springer, New York, (1971).
- [6] V. B. Osegove, Some extremal properties of generalized Appell polynomials, Soviet Math. Five, pp. 1651-1653, (1964).
- [7] E. D. Rainville, Special Functions, the Macmillan Company, New York, (1960).
- [8] I. M. Sheffer, Some properties of polynomial sets of type zero, Duke Math. J. 5, pp. 590-622, (1939).
- [9] J. F. Steffensen, The Poweroid, an extension of the mathematical notion of power, Acta Math. 73, pp. 333-366, (1941).

A. K. Shukla

Department of Mathematics
S. V. National Institute of Technology
Surat-395 007
India
e-mail : ajayshukla2@rediffmail.com,

and

S. J. Rapeli

Department of Mathematics

S. V. National Institute of Technology

Surat-395 007

India

e-mail : shrinu0711@gmail.com