

Proyecciones Journal of Mathematics
Vol. 32, N° 4, pp. 321-332, December 2013.
Universidad Católica del Norte
Antofagasta - Chile

Multiplication and Composition operators on $w_p(f)$

Kuldip Raj, Sunil K. Sharma and Seema Jamwal
Shri Mata Vaishno Devi University, India
Received : June 2012. Accepted : October 2013

Abstract

In this paper we characterize the boundedness, closed range, invertibility of the multiplication operators acting on sequence spaces $w_p(f)$ defined by a modulus function. We also make an efforts to study some properties of composition operators on these spaces.

Subclass [2000] : *Primary 47B20, Secondary 47B38.*

Keywords : *Modulus function, multiplication operator, composition operator, closed range, invertibility.*

1. Introduction and Preliminaries

A modulus function is a function $f : [0, \infty) \rightarrow [0, \infty)$ such that

1. $f(x) = 0$ if and only if $x = 0$;
2. $f(x + y) \leq f(x) + f(y)$ for all $x \geq 0, y \geq 0$;
3. f is increasing;
4. f is continuous from right at 0.

It follows that f must be continuous everywhere on $[0, \infty)$. The modulus function may be bounded or unbounded. For example, if we take $f(x) = \frac{x}{x+1}$, then $f(x)$ is bounded. If $f(x) = x^p$, $0 < p < 1$, then the modulus $f(x)$ is unbounded. Subsequently, modulus function has been discussed in ([2], [7], [9]) and many others.

For any sequence x , write

$$d_{mn} = d_{mn}(x) = \frac{1}{m+1} \sum_{i=0}^m x_{n+i}.$$

G. G. Lorentz [6] proved that

$$\hat{c} = \left\{ x : \lim_{m \rightarrow \infty} d_{mn}(x) \text{ exists uniformly in } n \right\}.$$

Khan M. A. [3] extend the definition of d_{mn} to $m = -1$ by taking $d_{-1,n} = x_{n-1}$, then write for $m, n \geq 0$

$$t_{mn} = t_{mn}(x) = d_{mn}(x) - d_{m-1,n}(x).$$

A straight forward calculation then show that

$$t_{mn} = \frac{1}{m(m+1)} \sum_{v=1}^m v(x_{n+v} - x_{n+v-1}).$$

If f is a modulus function and homogeneous of degree 1, then we define a sequence space as

$$w_p(f) = \left\{ x : \sup_n \sum_m m^{p-1} f(|t_{mn}(x)|^p) < \infty \right\}.$$

The space $w_p(f)$ with the norm

$$\|x\| = \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x)|^p) \right\}^{1/p}, \quad p \geq 1 \text{ for } x \in w_p(f)$$

is a Banach space.

Let $v : \mathbf{N} \rightarrow \mathbf{N}$ and $u : \mathbf{N} \rightarrow \mathbf{C}$ be two mappings.

Then the bounded linear transformations

$$T_v : w_p(f) \rightarrow w_p(f)$$

and

$$M_u : w_p(f) \rightarrow w_p(f)$$

defined by $(T_v h)(x) = h(v(x))$ and $(M_u h)(x) = u(x)h(x)$ are called composition and multiplication operators respectively. By $B(w_p(f))$, we denote the set of all bounded linear operators from $w_p(f)$ into itself and $[z(u)]$ denote, the set $\{n \in \mathbf{N} : u(n) = 0\}$. For more details about the study of multiplication and composition operators see ([1], [4], [5], [8], [10], [11]).

In this paper we study multiplication and composition operators acting on sequence spaces $w_p(f)$ defined by a modulus function.

2. Multiplication operators acting on sequence spaces defined by a modulus function

In this section we characterize multiplication operators acting on $w_p(f)$.

Theorem 2.1. *Let $M_u : w_p(f) \rightarrow w_p(f)$ be a linear transformation. Then M_u is a bounded operator if and only if there exists $M > 0$ such that*

$$f(|u(m)t_{mn}(x)|^p) \leq M f(|t_{mn}(x)|^p)$$

for all $m \in \mathbf{N}$.

Proof. Suppose that the condition of the theorem is true. For $x \in w_p(f)$, we have

$$\sup_n \sum_{m=1}^{\infty} m^{p-1} f(|u(m)t_{mn}(x)|^p) \leq M \sup_n \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x)|^p) < \infty.$$

Thus $M_u x \in w_p(f)$. Further,

$$\begin{aligned} \|M_u(x)\| &= \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|u(m)t_{mn}(x)|^p) \right\}^{1/p} \\ &\leq M \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x)|^p) \right\}^{1/p} \\ &\leq M \|x\|. \end{aligned}$$

This proves the continuity of M_u at the origin and hence everywhere in view of linearity of M_u .

Conversely, if the condition of the theorem were false, then for every integer $k > 0$ there exists $n_k \in N$ and $y_k = y \in \mathbf{R}^+$ such that

$$f(|u(n_k)t_{mn}(y_k)|^p) > kf(|t_{mn}(y_k)|^p)$$

Let $g_k = t_{mn}(y_k)\chi_{\{n_k\}}$. Then

$$\begin{aligned} k\|g_k\| &= k \left\| t_{mn}(y_k)\chi_{\{n_k\}} \right\| \\ &= k \left(\sup_n \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(y_k)|^p) \right)^{1/p} \\ &< \left(\sup_n \sum_{m=1}^{\infty} m^{p-1} f(|u(n_k)t_{mn}(y_k)|^p) \right)^{1/p} \\ &= \left(\sup_n \sum_{m=1}^{\infty} m^{p-1} f|M_u g_k|^p \right)^{1/p} \\ &= \|M_u g_k\|. \end{aligned}$$

This proves that M_u is not bounded. Hence the condition must be true.

Theorem 2.2. *Let $AM_u = M_uA$. Then A is a multiplication operator.*

Proof. Let $V = Ae$. Then

$$Ae_n = AM_{e_n}e = M_{e_n}Ae = M_{e_n}V = e_nV = Ve_n = M_Ve_n.$$

We now prove that V induces a multiplication operator. If V does not induce a bounded operator, then for every $k \in \mathbf{N}$, there exists $n_k \in \mathbf{N}$ such that

$$f\left(|V(n_k)t_{mn}(y_k)|^p\right) > mf\left(|t_{mn}(y_k)|^p\right).$$

Let $g_k = t_{mn}(y_k)e_{n_k}$. Then

$$\begin{aligned} k\|g_k\| &= k\|t_{mn}(y_k)e_{n_k}\| \\ &= k\left(\sup_n \sum_{m=1}^{\infty} m^{p-1}f\left(|t_{mn}(y_k)|\right)^p\right)^{1/p} \\ &< \left(\sup_n \sum_{m=1}^{\infty} m^{p-1}f\left(|V(n_k)t_{mn}(y_k)|\right)^p\right)^{1/p} \\ &= \left(\sup_n \sum_{m=1}^{\infty} m^{p-1}f|Ag_k|^p\right)^{1/p} \\ &= \|Ag_k\|, \end{aligned}$$

which contradicts the continuity of A . Hence A must be a bounded operator and $A = M_V$.

Theorem 2.3. *Let $M_u \in B(w_p(f))$. Then M_u is invertible if and only if there exists $\epsilon > 0$ such that*

$$f\left(|u(k)t_{mn}(y)|^p\right) \geq \epsilon f\left(|t_{mn}(y)|^p\right), \quad \forall p \in \mathbf{N} \text{ and } y \in \mathbf{R}^+.$$

Proof. We first assume that there exists $\epsilon > 0$ such that

$$f\left(|u(k)t_{mn}(y)|^p\right) \geq \epsilon f\left(|t_{mn}(y)|^p\right), \quad \forall p \in \mathbf{N} \text{ and } y \in \mathbf{R}^+.$$

Now

$$\begin{aligned} \epsilon f \left[\frac{|t_{mn}(y)|^p}{|u(k)|^p} \right] &\leq f \left[|u(k)|^p \cdot \frac{|t_{mn}(y)|^p}{|u(k)|^p} \right] \\ &= f \left(-|t_{mn}(y)|^p \right) \text{ or} \\ f \left[\frac{1}{|u(k)|^p} |t_{mn}(y)|^p \right] &\leq \frac{1}{\epsilon} f \left(|t_{mn}(y)|^p \right), \quad \forall p \in \mathbf{N}. \end{aligned}$$

This proves that M_V is a bounded operator, where $V = \frac{1}{u}$. Clearly M_V is inverse of M_u .

Conversely, suppose that M_u is invertible with M_V as its inverse. Clearly $V = \frac{1}{u}$. Hence by continuity of M_V , there exists $M > 0$ such that

$$f \left(|V(k)t_{mn}(y)|^p \right) \leq M f \left(|t_{mn}(y)|^p \right), \quad \forall k \in \mathbf{N} \text{ and } y \in \mathbf{R}^+.$$

Or equivalently

$$f \left[\frac{1}{|u(k)|} |t_{mn}(y)|^p \right] \leq M f \left(|t_{mn}(y)|^p \right).$$

Taking

$$|t_{mn}(y)| = |u(k)t_{mn}(y)|,$$

we get

$$f \left(|t_{mn}(y)|^p \right) \leq M f \left(|u(k)t_{mn}(y)|^p \right)$$

or

$$f \left(|u(k)t_{mn}(y)|^p \right) \geq \frac{1}{M} f \left(|t_{mn}(y)|^p \right) \quad \forall k \in \mathbf{N}.$$

Taking $\epsilon = \frac{1}{M}$, we get

$$f \left(|u(k)t_{mn}(y)|^p \right) \geq \epsilon f \left(|t_{mn}(y)|^p \right).$$

Hence the condition must be true.

Theorem 2.4. *Let $M_u \in B(w_p(f))$. Then M_u is Fredholm if and only if*

- (i) $[Z(u)]$ is a finite set
- (ii) there exists $\epsilon > 0$ such that

$$f\left(|u(k)t_{mn}(y)|^p\right) \geq \epsilon f\left(|t_{mn}(x)|^p\right) \quad \forall m \in [Z(u)]'$$

Proof. If $[Z(u)]$ is a finite set, then $\ker M_u$ is finite dimensional. From the condition (ii), M_u has closed range.

Moreover $\dim(w_p(f)/\text{ran}M_u)$ is finite. This proves that M_u is Fredholm.

The converse of the theorem is obvious.

Corollary 2.5. *Let $M_u \in B(w_p(f))$. Then M_u has closed range if and only if there exists $\delta > 0$ such that*

$$f\left(|u(k)t_{mn}(y)|^p\right) \geq \delta f\left(|t_{mn}(y)|^p\right), \quad \forall k \in [Z(u)]' \text{ and } y \in \mathbf{R}^+.$$

Proof. Assume that the condition of the theorem is true. Let $h \in \overline{\text{ran}M_u}$.

Then there exists a sequence $\{h_n\}$ such that $M_u h_n \rightarrow h$ that is $\|M_u h_n - M_u h\| \rightarrow 0$ as $n \rightarrow \infty$. Now $\{M_u h_n\}$ is a Cauchy sequence. Therefore for every $\epsilon > 0$ there exists $n_0 \in \mathbf{N}$ such that

$$\|M_u t_{mn} h_n - M_u t_{mn} h_k\| < \epsilon \quad \forall n, k \geq n_0.$$

Now

$$\begin{aligned} \delta \sup_{n \in [Z(u)]'} \sum_{m=1}^{\infty} m^{p-1} f\left(|t_{mn}(h_n - h_k)|^p\right) &\leq \sup_{n \in [Z(u)]'} \sum_{m=1}^{\infty} m^{p-1} f\left(|u(m)t_{mn}(h_n - h_k)|^p\right) \\ &< \epsilon \quad \forall n, k \geq n_0. \end{aligned} \tag{1}$$

Define

$$\tilde{h}_n(k) = \begin{cases} h_n(k), & \text{if } m \in [Z(u)]' \\ 0, & \text{elsewhere.} \end{cases}$$

Then from (1) it follows that $\{\tilde{h}_n\}$ is a Cauchy sequence in $w_p(f)$. But $w_p(f)$ is complete.

Therefore there exists $\tilde{h} \in w_p(f)$ such that $\tilde{h}_n \rightarrow \tilde{h}$. Hence by continuity of M_u , we get $M_u h_n = M_u \tilde{h}_n \rightarrow M_u \tilde{h}$. Hence $h = M_u \tilde{h}$ so that $h \in \text{ran} M_u$. Thus M_u has closed range.

Conversely, if the condition of the theorem were false, then for every positive integer k there exists $n_k \in N$ and $y_k \in \mathbf{R}^+$ such that

$$f(|u(n_k)t_{mn}(y_k)|^p) < 1/k f(|t_{mn}y_k|^p).$$

Let $g_k = t_{mn}y_k \chi_{\{n_k\}}$.

Then

$$\begin{aligned} \|M_u g_k\| &= \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|u \cdot g_k|^p) \right\}^{1/p} \\ &= \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|u(n_k)t_{mn}(y_k)|^p) \right\}^{1/p} \\ &\leq 1/k \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(y_k)|^p) \right\}^{1/p} \\ &= 1/k \|g_k\|. \end{aligned}$$

This proves that M_u is not bounded away from zero so that M_u does not have closed range.

3. Composition operators acting on sequence spaces defined by a modulus function

In this section we study some properties of composition operators on $w_p(f)$.

Theorem 3.1. *Let $T_v : w_p(f) \rightarrow w_p(f)$ be a linear transformation. Then T_v is a bounded operator if there exists $M > 0$ such that*

$$\sum_{k \in v^{-1}(n)} m^{p-1} f(|t_{mk}(x)|^p) \leq M m^{p-1} f(|t_{mn}(x)|^p).$$

Proof. Suppose that the condition of the theorem is true. If $x \in w_p(f)$, then

$$\sup_n \sum_{m=1}^{\infty} \sum_{k \in v^{-1}(n)} m^{p-1} f(|t_{mk}(x)|^p) \leq M \sup_n \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x)|^p)$$

$< \infty$, which shows that $T_v x \in w_p(f)$. Further,

$$\begin{aligned} \|T_v x\|_f &= \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x \circ v(k))|^p) \right\}^{1/p} \\ &= \sup_n \left\{ \sum_{m=1}^{\infty} \sum_{k \in v^{-1}(n)} m^{p-1} f(|t_{mk}x|^p) \right\}^{1/p} \\ &\leq M \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x)|^p) \right\}^{1/p} \\ &\leq M \|x\|_f. \end{aligned} \tag{2}$$

The continuity of T_v at origin follows from the inequality (2). Since T_v is linear, so it is continuous everywhere.

Theorem 3.2. Let $T_v \in B(w_p(f))$. Then T_v has closed range if there exists $\delta > 0$ such that

$$\sum_{k \in v^{-1}(n)} m^{p-1} f(|t_{mk}(x)|^p) \geq \delta m^{p-1} f(|t_{mn}(x)|^p) \text{ for every } m \in \mathbf{N}. \tag{3}$$

Proof. We assume that the condition (3) is true. We have to show that T_v has closed range. Let $x \in \overline{\text{ran} T_v}$ and let $\{x^i\}$ be a sequence in $w_p(f)$ such that $T_v x^i \rightarrow x$. Then for every $\epsilon > 0$ there exists positive integer n_0 such that

$$\|T_v x^i - T_v x^j\| < \epsilon \quad \forall i, j \geq n_0.$$

Equivalently,

$$\epsilon > \sup_n \left\{ \sum_{m=1}^{\infty} \sum_{k \in v^{-1}(n)} m^{p-1} f(|t_{mk}(x^i \circ v(k)) - x^j \circ v(k)|^p) \right\}^{1/p}$$

$$\begin{aligned}
&\geq \delta \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x^i - x^j)|^p) \right\}^{1/p} \\
&= \delta \|x^i - x^j\|, \quad \forall i, j \geq n_0(4)
\end{aligned}$$

from (4) it follows that $\{x^i\}$ is a Cauchy sequence in $w_p(f)$. In view of completeness of $w_p(f)$, there exists $y \in w_p(f)$ such that $x^i \rightarrow y$. From the continuity of T_v , $T_v x^i \rightarrow T_v y$. Hence $x = T_v y$ so that $x \in \text{ran} T_v$. Hence $\text{ran} T_v$ is closed.

Theorem 3.3. *Let $T_v \in B(w_p(f))$. Then T_v is an isometry if*

$$\sum_{k \in v^{-1}(n)} m^{p-1} f(|t_{mk}(x)|^p) = m^{p-1} f(|t_{mn}(x)|^p).$$

Proof. If the condition of the theorem is satisfied, then for every $x \in w_p(f)$, we have

$$\begin{aligned}
\|T_v x\| &= \sup_n \left\{ \sum_{m=1}^{\infty} \sum_{k \in v^{-1}(m)} m^{p-1} f(|t_{mk}x|^p) \right\}^{1/p} \\
&= \sup_n \left\{ \sum_{m=1}^{\infty} m^{p-1} f(|t_{mn}(x)|^p) \right\}^{1/p} \\
&= \|x\|.
\end{aligned}$$

Hence T_v is an isometry.

Theorem 3.4. *Let $T_v \in B(w_p(f))$. If T_v is an isometry, then*

$$\sup_n \sum_{k \in v^{-1}(m)} k^{p-1} f(|t_{nk}(x)|^p) = \sup_n m^{p-1} f(|t_{mn}x|^p).$$

Proof. The proof is trivial.

References

- [1] M. B. Abrahmese, Multiplication operators, Hilbert space operators. *Lecture notes in Mathematics*, 693 : pp. 17-36, (1978).
- [2] T. Bilgen, On statistical convergence. *An. Univ. Timisoara Ser. Math. Inform.*, 32 : pp. 3-7, (1994).
- [3] M. A. Khan, Some sequence spaces with an index defined by a modulus function. *Thai J. Math.*, 2 : 259-264, (2004).
- [4] B. S. Komal and Kuldip Raj, Multiplication operators induced by operator valued maps. *Int. J. Contemp. Math. Sci.*, Vol.3 : pp. 667-673, (2008).
- [5] B. S. Komal and P. S. Singh, Composition operators on the space of entire functions. *Kodai Math. J.*, 14 : pp. 463-469, (1991).
- [6] G. G. Lorentz, A contribution to the theory of divergent series. *Acta. Math.*, 80 : pp. 167-190, (1948).
- [7] E. Malkowsky and E. Savas, Some λ -sequence spaces defined by a modulus. *Arch. Math.*, 36 : pp. 219-228, (2000).
- [8] Kuldip Raj, B. S. Komal and Vinay Khosla, Composition operators on sequence spaces of entire functions. *Int. Electron. J. Pure Appl. Math.*, 1 : pp. 469-474, (2010).
- [9] E. Savas, On some generalized sequence spaces defined by a modulus. *Indian J. pure Appl. Math.*, 30 : pp. 459-464, (1999).
- [10] R. K. Singh and J. S. Manhas, Composition operators on function spaces. *North-Holland*, (1993).
- [11] H. Takagi and K. Yokouchi, Multiplication and composition operators between L^p -spaces. *Contemp. Math.*, 232 : pp. 321-338, (1999).

Kuldip Raj

School of Mathematics
Shri Mata Vaishno Devi University
Katra-182320, J&K,
India
e-mail : kuldipraj68@gmail.com

Sunil K. Sharma

School of Mathematics
Shri Mata Vaishno Devi University
Katra-182320, J&K,
India
e-mail : sunilksharma42@yahoo.co.in

and

Seema Jamwal

School of Mathematics
Shri Mata Vaishno Devi University
Katra-182320, J&K,
India
e-mail : seemajamwal8@gmail.com