

# Population density and reproduction of two Peruvian leaf-eared mice (*Phyllotis* spp.)

Densidad poblacional y reproducción de dos ratones orejados del Perú (*Phyllotis* spp.)

MARGARITA ARANA, OSWALDO RAMÍREZ<sup>1</sup>, SERGIO SANTA MARÍA, CESAR KUNIMOTO, RODRIGO VELARDE,  
CECILIA DE LA CRUZ & MARÍA LUISA RUÍZ

Departamento de Ciencias Biológicas y Fisiológicas, Facultad de Ciencias, Universidad Peruana  
Cayetano Heredia, Avenida Honorio Delgado 430, Urbanización Ingeniería,  
San Martín de Porres, Lima, Perú; e-mail<sup>1</sup>: oramirez@upch.edu.pe

## ABSTRACT

Population density and reproduction were studied in two leaf-eared mice (*Phyllotis* spp.) in the Peruvian locality known as Lomas de Lachay. Density varied between 0-3 individuals ha<sup>-1</sup> in *P. andium*, and between 0-12 individuals ha<sup>-1</sup> in *P. amicus*. An annual cycle of population abundance was observed in both species. *Phyllotis amicus* showed no seasonal reproductive activity, but *P. andium* had seasonal reproduction with pregnant females principally observed between July and September. Differences in juvenile weight were related to litter size. It is likely that two young was the optimal litter size in the harsh and seasonal xeric environment of Lomas de Lachay. Our results suggest that *P. amicus* maintains low, highly dispersed populations in a harsh environment, and can reproduce at any time of the year, whereas *P. andium* appears to be a more irruptive species closely tied to rainfall patterns.

**Key words:** ecology, *Phyllotis*, rodents, population, desert.

## RESUMEN

La densidad poblacional y reproducción de dos ratones orejados (*Phyllotis* spp.) fueron estudiadas en la localidad peruana Lomas de Lachay. La densidad varió entre 0-3 individuos ha<sup>-1</sup> en *P. andium*, y entre 0-12 individuos ha<sup>-1</sup> en *P. amicus*. En ambas especies se observaron cambios poblacionales que sugieren la existencia de ciclos anuales en la abundancia poblacional. El patrón reproductivo en las dos especies de *Phyllotis* fue claramente distinto. *Phyllotis amicus* mostró una actividad reproductiva no estacional, mientras que *P. andium* tuvo una reproducción estacional observándose hembras preñadas principalmente entre julio y septiembre. La diferencia en el peso de los juveniles estuvo relacionada con el tamaño de la camada. Es probable que en las Lomas de Lachay el tamaño de camada óptimo sea de dos individuos. Nuestros resultados sugieren que *P. amicus* es una especie que mantiene poblaciones bajas, altamente dispersas en áreas desérticas, y que puede reproducirse durante todo el año; mientras que *P. andium* parece ser una especie que irrumpe más en ambientes arbustivos dependientes de patrones de precipitación pluvial.

**Palabras clave:** ecología, *Phyllotis*, roedores, población, desierto.

## INTRODUCTION

Leaf-eared mice (genus *Phyllotis*, Sigmodontinae) are one of the most abundant small mammals in the Peruvian Andes. Eight of 13 species inhabit this area, where a predominance of xeric habitats exists with sparse vegetation (Pearson 1958, Steppan 1995, 1998). Ecological studies of the Peruvian species of *Phyllotis* are scarce and reproduction data have been only occasionally reported. Two species of leaf-eared mice (*P. amicus* and *P. andium*) can be found in a unique seasonal habitat, the Lomas of Lachay, located on the central coast of Peru, 105 km north of Lima. Here, we present information on population

density, movements, and reproduction (in natural and captive conditions) for these mice species.

## MATERIAL AND METHODS

The fieldwork was done in the Teatinos pass, located on the western side of the Lachay National Reserve (11°21'00"-11°21'58" S, 77°21'18"-77°22'25" W). Mean annual precipitation in the last four years (1998-2001) was 0.88, 0.29, 0.41, and 0.42 mm, respectively. One 1-ha grid (altitude: 420 m) with 81 trap stations (9 x 9, 10-m intervals, one trap per station) was censused monthly from June 2000 to December 2001. A second (altitude:

350 m), similarly-sized grid was censused between November 2000 and December 2001. Baited Sherman traps were placed at stations immediately before sunset and checked the following morning for two consecutive days each month. Animals were marked by toe clipping and released at the point of capture. Sex, reproductive condition, weight, and grid station were recorded. Traps were closed during the daytime. Population abundance was determined by the capture-mark-recapture method using the Lincoln-Petersen index as modified by Chapman (1951). In order to study movements, we analyzed the location of consecutive captures of individuals through time, and the maximum time between first and last capture. Movement distance (minimum distance between the two furthest points of capture) also was calculated.

Reproductive activity of both *Phyllotis* species was assessed from examination of captured animals inside and outside grids. In addition, individuals of both species of *Phyllotis* (captured outside grids) were kept and bred in captivity. Each animal was housed in small cages 30 x 18 x 13 cm using wood shavings and paper as bedding materials. Animals were fed with fresh carrots, lettuce, sweet potato, sprouted wheat, and corn. Water was not provided to the animals after the first months as they never drank. Animals were maintained at an average temperature of  $23 \pm 5$  ( $\pm$  SD) °C. For reproduction, one male was placed in a large cage (36 x 32 x 20 cm) for at least two days before introduction of the female. Pairs were maintained together for two weeks, after which the females were returned to small cages whereas the males were maintained in the large cages until a new mate was introduced. Animals were checked daily to record births and deaths in the colony. To analyze the effect of litter size on the weight of young, we recorded body weight each 10-20 days for 90 or more days after birth. Regression analyses between weight versus age were conducted for each litter size. Homogeneity of slopes were tested, and analyses of covariance were conducted to test for statistical differences ( $P < 0.05$ ) among the regressions.

## RESULTS

Population densities of both *Phyllotis* species are shown in Fig. 1. Very small numbers of individuals per ha were observed during almost the entire study. Only *P. amicus* in the low grid showed a higher population density (12.3 individuals ha<sup>-1</sup>) between May and July of 2001. *Phyllotis andium* was more abundant in the high grid than in the low

grid, and exhibited the highest numbers between July and August of 2000. Six *P. amicus* individuals were recaptured for more than three months and one for eight months. Using data from these individuals, movement distances varied between 22.8 and 78.8 m ( $\bar{X} = 44.5 \pm 17.9$  m). Only one *P. andium* individual was recaptured one month later. Using the three *P. andium* individuals captured at least two times in the same month, the movement distance between captures varied between 40.8 and 68.8 m ( $\bar{X} = 50.5 \pm 12.9$  m).

In captivity, 76 and 12 matings were conducted between individuals of *P. andium* and *P. amicus*, respectively. Twenty litters with a total of 45 young were obtained in *P. andium*, whereas four litters with eight young were obtained in *P. amicus*. The gestation time in both species was 24 days. Litter size varied between one and four young in *P. andium* and between one and three pups in *Phyllotis amicus*. Fig. 2 shows the number of pregnant females captured in Lomas de Lachay




Fig. 1: Variation in the population density of *Phyllotis andium* and *P. amicus*. For each species two grids were examined: (A) June 2000 – December 2001, and (B) November 2000 – December 2001.

Variación en la densidad poblacional de *Phyllotis andium* y *P. amicus*. Se estudiaron dos cuadrantes para cada especie: (A) Junio 2000 – diciembre 2001, y (B) noviembre 2000 – diciembre 2001.




Fig. 2: Number of pregnant females captured in Lachay lomas by month. The black portion of the bars show the number of pregnant females and the white portion shows the number of non pregnant females.

Número de hembras preñadas capturadas mensualmente en las lomas de Lachay. La parte negra de las barras indica el número de hembras preñadas y la parte blanca muestra el número de hembras no preñadas.

each month. *Phyllotis amicus* showed no seasonal reproduction, with pregnant females throughout the year. In contrast, *P. andium* showed clear seasonal reproductive activity between July and September. In the laboratory, 35 % of female *P. andium* that became pregnant had litters with two young, whereas 30, 20 and 15 % had litters of one, four and three young, respectively. An ANCOVA showed significant differences in the regressions between weight and age with different litter size (Fig. 3), indicating a relationship between weight and litter size. Heavier animals were from litters of one young whereas smaller animals came from litters with four young. Young (irrespective of litter size) were weaned 30 days after birth. Mean weight of young at weaning was  $14.4 \pm 2.6$  g. Young reached mean weights of  $17 \pm 3.4$  and  $20.8 \pm 4.7$  g during their second and third months of life, respectively. In captivity, females less than seven months of age were not reproductively active.

DISCUSSION

This is the first report on the population ecology of two species of leaf-eared mice in the Lomas de Lachay, and the second that comes from Peruvian lomas (Péfaur et al. 1979, Pearson 1975). Pearson & Ralph (1978) monitored *Phyllotis limatus* in a 2.02-ha grid in a loma near Tacna, 1,200 km south of Lima. They found that the population density

TABLE 1

Movement distances and length of residence of *Phyllotis amicus* and *Phyllotis andium*

Distancia de desplazamiento y tiempo de residencia de *Phyllotis amicus* y *Phyllotis andium*

Species	Sex	Movement distance (m)	Number of captures	Length of residence (days)
<i>Phyllotis amicus</i>	♀	29.60	5	245
	♀	22.80	6	91
	♀	43.20	4	122
	♂	50.40	5	127
	♂	78.80	6	157
	♂	42.00	3	100
<i>Phyllotis andium</i>	♀	40.80	4	37
	♂	68.00	3	3
	♂	42.00	3	3




Fig. 3: Regression lines between weight and age for different litter size in *P. andium*.

Regresión lineal entre peso y edad en camadas de diferentes tamaños de *P. andium*.

was 6.5 individuals  $ha^{-1}$ , but considered this value to be an overestimate because the study occurred in November when the breeding season was ending and many young *Phyllotis* were in the population. After excluding these young, the population density was 3.9 individuals  $ha^{-1}$ . Our results with *P. andium* and *P. amicus* showed similar values of density and suggest an annual cycle (Fig. 1), a pattern that has been previously reported for several South American rodents (Crespo et al. 1970, González et al. 1982, Murúa 1983, Murúa & González 1985, Murúa & González 1986, Murúa et al. 1986). Higher population densities were observed during the wet season (June–August) in both species. One interesting feature was the absence of individuals of *P. andium* during most of the study period in both grids. When traps were placed in other areas of Lomas de Lachay, the result for *P. andium* followed the same trend observed in the grids. Therefore, only occasional individuals appear to be present in the lomas except between July and August when their numbers increase. Captures of *P. amicus* outside of the grids were very low, and the individuals trapped were always in the most xeric and sparse habitats at low elevations in Lomas de Lachay. This suggests some spatial segregation between these two species of *Phyllotis*. *Phyllotis andium*

occupies most of the area when in low numbers whereas *P. amicus* occurs in more restricted, xeric areas in the lowest parts of Lomas de Lachay. This segregation agrees with findings of Pizzimenti & De Salle (1980) who also captured *P. amicus* isolated from its congeners. Both species of *Phyllotis* had nocturnal activity, and showed an aggregated distribution throughout the year suggesting a rather social type of spatial organization, which could be the result of some limiting factors such as substrate or shelter.

Mean distances traveled were 44.5 and 50.5 m for *P. amicus* and *P. andium*, respectively. Pearson & Ralph (1978) found a movement distance of 36 m for *P. limatus* at Morro Sama, a loma in southern Peru. Although this value is within the range of distances recorded in *P. amicus* (22.8–78.8 m) and close to the minimum value found in *P. andium* (40.8 and 68.8 m), it is likely an underestimate due to the grid being trapped for only six nights in Morro Sama. Nevertheless, these results suggest that *Phyllotis* spp. restrict their movements in lomas-type ecosystems.

Reproductive activity differed between the two species of *Phyllotis* (Fig. 2). Pregnant females of *P. andium* were found mainly between July and September, whereas pregnant females of *P. amicus* were found throughout the year. These results

suggest that *P. andium* reproduce seasonally and that breeding coincides with the wet period (winter; with precipitations above to 1.0 mm monthly). *Phyllotis amicus* lacked seasonal reproductive activity. A similar result was found by Williams (1990), who captured two pregnant *P. amicus* females in Lomas de Lachay during May and September. Differences between the reproductive patterns of *P. andium* and *P. amicus* may be related to their food habits. Although both species have broadly omnivorous feeding habits, *P. amicus* eats more insects than any other species of *Phyllotis* (Pizzimenti & De Salle 1980), and therefore, this species may have a reproductive activity with a lesser dependence on vegetation. This finding is relevant due to the higher population density of *P. amicus* during May and July, which does not seem to be due to juvenile recruitment. One untested hypothesis is that mobility increases in the population, as it had been reported in other sigmodontine rodents (Murúa & González 1986).

Differences in the weight of juveniles born in captivity are related to the litter size as shown in Fig. 3. Young from a litter of 1-2 individuals reach a higher weight in less time, thus likely increasing their survival rate. This suggests that there is a balance between the number and survivorship of young (as measured by growth and weight). Since the majority of pregnant *P. andium* females had a litter size of two, such figure could be the optimal litter size under the regular climatic conditions at Lomas de Lachay. However, during anomalous climatic conditions when food resources are over abundant (i.e., after El Niño Southern Oscillations, ENSO events), outbreaks of some *Phyllotis* species may occur (Pearson 1975, Jaksic et al. 1977, Péfaur et al. 1979, Meserve et al. 1995, Lima et al. 1999a, 1999b). Meserve et al. (1995) reported a dramatic increase in the population size of *P. darwini* in north-central Chile as a consequence of the 1991-1992 ENSO event. The population size of this species never exceeded 9 individuals  $0.56 \text{ ha}^{-1}$  between 1989 and 1991, but reached a maximum of 25 in February 1992 and of 39 in May 1993. On the other hand, *P. darwini* reached 200 individuals  $\text{ha}^{-1}$  during summer 1988 in Aucó, north-central Chile (Lima et al. 1999a, 1999b). Given that *P. darwini* is capable of producing many litters within a short period of time (Meserve & Le Boulengé, 1987), and that the litter size in this species varies between 4.2 (Walker et al. 1984) and 5.2 (Fulk 1975, Meserve & Le Boulengé 1987), the rapid increase of population size during an outbreak ("ratada") indicates that *Phyllotis* spp. have a reproductive potential largely constrained by environmental factors. Therefore, in a xeric, harsh, seasonal

environment such as lomas ecosystems, a small litter size of two provides greater survival potential than larger litter sizes (4-6).

Our results showed no reproductive activity of females younger than seven months in *P. andium*. This may explain the relatively delayed response in the population changes of *Phyllotis* as a biological effect of ENSO events (Pearson 1975, Péfaur et al. 1979, Meserve et al. 1995). Other recent reports have suggested an interaction between delayed density-dependent factors and rainfall at demographic and population dynamic levels as the key element for understanding outbreaks of *P. darwini* in semiarid regions of Chile (Lima et al. 1999a, 1999b, Lima & Jaksic 1999a, 1999b). Our short-term study did not allow us to evaluate this hypothesis in the species here studied.

The apparent year-around reproduction of *P. amicus* (in contrast to *P. andium*) and its very low population level may be a strategy that enables this species to persist in the lomas regardless of environmental conditions. Indeed, our results suggest that *P. amicus* is a species that maintains low, highly dispersed populations in a harsh environment, and can reproduce at any time of year, whereas *P. andium* appears to be a more irruptive species closely tied to rainfall patterns.

#### ACKNOWLEDGMENTS

This research was supported in part by a grant from the Red Impacto Biológico El Niño (RIBEN) / Concejo Nacional de Ciencia y Tecnología (CONCYTEC). We thank to David Velarde, Chief of Lachay National Reserve and his personal for giving us logistic facilities during the field work and precipitation data. We are very grateful to Peter Meserve for his comments and suggestions on this article.

#### LITERATURE CITED

- CHAPMAN DG (1951) Some properties of the hypergeometric distribution with applications to zoological sample censuses. University of California Publications Statistics 1: 131-160.
- CRESPO JA, MS SABATTINI, MJ PIANTANIDA & G DE VILLAFANE (1970) Estudios ecológicos sobre roedores silvestres. Observaciones sobre densidad, reproducción y estructura comunitaria de roedores silvestres en el sur de Córdoba. Comisión Nacional de Estudio y Lucha contra la Fiebre Hemorrágica Argentina, Ministerio de Bienestar Social, Secretaría de Salud Pública, Buenos Aires, Argentina. 45 pp.

- FULK GW (1975) Population ecology of rodents in the semiarid shrublands of Chile. Occasional Papers, The Museum, Texas Tech University 33: 1-40.
- GONZÁLEZ LA, R MURÚA & R FEITO (1982) Densidad poblacional y padrones de actividad espacial de *Akodon olivaceus* (Rodentia, Cricetidae) en hábitats diferentes. In: Salinas P (ed) Actas de Séptimo Congreso Latinoamericano de Zoología, Zoología Neotropical: 935-947. Mérida, Venezuela.
- JAKSIC FM, SI SILVA, PL MESERVE & JR GUTIÉRREZ (1997) A long-term study of vertebrate predator responses to an El Niño (ENSO) disturbance in western South America. *Oikos* 78: 341-354.
- LIMA M & FM JAKSIC (1999a) Population rate of change in the leaf-eared mouse: the role of density-dependence, seasonality and rainfall. *Australian Journal of Ecology* 24: 110-116.
- LIMA M & FM JAKSIC (1999b) Population dynamics of three Neotropical small mammals: time series models and the role of delayed density-dependence in population irruptions. *Australian Journal of Ecology* 24: 25-34.
- LIMA M, JE KEYMER & FM JAKSIC (1999a) El Niño-Southern Oscillation-driven rainfall variability and delayed dependence cause rodent outbreaks in western South America: linking demography and population dynamics. *American Naturalist* 153: 476-491.
- LIMA M, PA MARQUET & FM JAKSIC (1999b) El Niño events, precipitation patterns, and rodent outbreaks are statistically associated in semiarid Chile. *Ecography* 22: 213-218.
- MESERVE PL & ER LE BOULENGÉ (1987) Population dynamics and ecology of small mammals in the northern Chilean semiarid region. In: Patterson BD & RM Timm (eds) Studies in Neotropical mammalogy: essays in honor of Philip Hershkovitz. *Fieldiana: Zoology* 39: 413-431.
- MESERVE PL, JA YUNGER, JR GUTIÉRREZ, LC CONTRERAS, WB MILSTEAD, BK LANG, KL CRAMER, S HERRERA, VO LAGOS, SI SILVA, EL TABILO, MA TORREALBA & FM JAKSIC (1995) Heterogeneous responses of small mammals to an El Niño Southern Oscillation event in northcentral semiarid Chile and the importance of ecological scale. *Journal of Mammalogy* 76: 580-595.
- MURÚA R (1983) Dinámica de los números en *Akodon olivaceus brachiotis* (Rodentia: Cricetidae). *Medio Ambiente (Chile)* 6: 12-18.
- MURÚA R & LA GONZÁLEZ (1985) A cycling population of *Akodon olivaceus* (Cricetidae) in a temperate rain forest in Chile. *Acta Zoológica Fennica* 173: 77-79.
- MURÚA R & LA GONZÁLEZ (1986) Regulation of number in two Neotropical rodent species in southern Chile. *Revista Chilena de Historia Natural* 59: 193-200.
- MURÚA R, LA GONZÁLEZ & PL MESERVE (1986) Population ecology of *Oryzomys longicaudatus philippii* (Rodentia: Cricetidae) in southern Chile. *Journal of Animal Ecology* 55: 281-293.
- PEARSON OP (1975) An outbreak of mice in the coastal desert of Peru. *Mammalia* 39: 375-386.
- PEARSON OP (1958) A taxonomic revision of the rodent genus *Phyllotis*. University of California Publications in Zoology 56: 391-496.
- PEARSON OP & CP RALPH (1978) The diversity and abundance of vertebrates along an altitudinal gradient in Peru. *Memorias del Museo de Historia Natural "Javier Prado"* 18: 1-97.
- PÉFAUR JE, JL YÁÑEZ & FM JAKSIC (1979) Biological and environmental aspects of a mouse outbreak in the semi-arid region of Chile. *Mammalia* 43: 313-322.
- PIZZIMENTI JJ & R DE SALLE (1980) Dietary and morphometric variation in some Peruvian rodent communities: the effect of feeding strategy on evolution. *Biological Journal of the Linnean Society* 13: 263-285.
- STEPHAN S (1995) Revision of the tribe Phyllotini (Rodentia: Sigmodontinae), with a phylogenetic hypothesis for the Sigmodontinae. *Fieldiana: Zoology* 80: 1-112.
- STEPHAN S (1998) Phylogenetic relationships and species limits within *Phyllotis* (Rodentia: Sigmodontinae): concordance between mtDNA sequence and morphology. *Journal of Mammalogy* 79: 573-593.
- WALKER LI, AE SPOTORNO & J ARRAU (1984) Cytogenetic and reproductive studies of two nominal subspecies of *Phyllotis darwini* and their experimental hybrids. *Journal of Mammalogy* 65: 220-230.
- WILLIAMS M (1990) Notas sobre reproducción de *Phyllotis amicus* Thomas (Rodentia: Cricetidae) en el campo y laboratorio. *Biota (Perú)* 96: 9-17.

Associate Editor: F. Jaksic

Received March 26, 2002; accepted August 21, 2002